arXiv:math/0605297v1 [math.AG] 11 May 2006

L_∞-ALGEBRAS, CARTAN HOMOTOPIES AND PERIOD MAPS

DOMENICO FIORENZA AND MARCO MANETTI

Abstract. We prove that, for every compact Kähler manifold, the period map of its Kuranishi family is induced by a natural L_∞-morphism. This implies, by standard facts about L_∞-algebras, that the period map is a "morphism of deformation theories" and then commutes with all deformation theoretic constructions (e.g. obstructions).

Contents

Introduction 2

Part 1. L_∞ structures on mapping cones 3
1. Conventions on graded vector spaces 4
2. Differential graded Lie algebras and L_∞-algebras 5
3. The suspended mapping cone of $\chi\colon L \to M$. 7
4. The L_∞ structure on C_χ 8
5. The functors MC_χ and Def_χ revisited 11
6. Functoriality 13
7. Cartan homotopies 14
Appendix A: gauge vs. homotopy 15

Part 2. dg-Grassmann functors 16
8. The coarse dg-Grassmannian 17
9. Infinitesimal study 18
10. Homotopy invariance 21

Part 3. The universal period map 24
11. The Kodaira-Spencer DGLA 24
12. The period map 27
13. Trasversality 29
References 30

Date: May 9, 2006.

1991 *Mathematics Subject Classification.* 14D07, 17B70, 13D10.

Key words and phrases. Differential graded Lie algebras, symmetric coalgebras, L_∞-algebras, functors of Artin rings, Kähler manifolds, period map.

INTRODUCTION

The homotopy Lie algebra approach to deformation theory over a field $\mathbb{K}$ of characteristic 0 is based on two creeds.
First creed: let $\mathcal{M}$ be a moduli space (for some classification problem) and x a point of $\mathcal{M}$; then the geometry of the formal neighbourhood of x in $\mathcal{M}$ is encoded into an (homotopy class of) L_∞-algebra(s). More precisely, there exists an L_∞-algebra $\mathfrak{g}$, defined up to quasiisomorphism, such that, for any local Artinian $\mathbb{K}$-algebras B with maximal ideal $\mathfrak{m}_B$ and residue field $\mathbb{K}$ one has

$$\{\phi\colon \operatorname{Spec}(B) \to \mathcal{M} \mid \phi(\operatorname{Spec}(\mathbb{K})) = x\} = \operatorname{Def}_{\mathfrak{g}}(B),$$

where

$$\operatorname{Def}_{\mathfrak{g}}(B) = \frac{\{\text{Solutions of the Maurer-Cartan equation in } \mathfrak{g}\otimes_{\mathbb{K}}\mathfrak{m}_B\}}{\text{homotopy equivalence}}.$$

Second creed: every "natural" morphism between formal pointed moduli spaces is induced by an L_∞-morphism between the associated L_∞-algebras.
In this paper we shall explicate the first creed for Grassmannians using a general construction that also applies to other moduli spaces (dg-Grassmannians, Quot and Hilbert schemes, Brill-Noether loci etc.) and the second creed for the universal period map, intended as a natural morphism of moduli spaces: from deformations of a compact Kähler manifold to deformations of the Hodge filtration of its De Rham cohomology.

More concretely, let's denote by

$$\mathcal{X} \to (S,0), \qquad \mathcal{X} = \bigcup_{t\in S} X_t,$$

the Kuranishi family of a compact Kähler manifold $X = X_0$; let $p \geq 0$ be a fixed integer and consider the period map [33, 10.1.2]

$$\mathcal{P}^p\colon (S,0) \to \operatorname{Grass}(H^*(X,\mathbb{C})) = \prod_i \operatorname{Grass}(H^i(X,\mathbb{C})),$$

$$t \mapsto F^pH^*(X_t,\mathbb{C}) = \prod_i F^pH^i(X_t,\mathbb{C}).$$

Griffiths proved [11] that $\mathcal{P}^p$ is a holomorphic map and its differential $d\mathcal{P}^p$ is the same of the contraction map

$$\boldsymbol{i}\colon H^1(X,T_X) \to \bigoplus_i \operatorname{Hom}\left(F^pH^i(X,\mathbb{C}), \frac{H^i(X,\mathbb{C})}{F^pH^i(X,\mathbb{C})}\right), \qquad \boldsymbol{i}_\xi(\omega) = \xi \lrcorner \omega.$$

It is also known [3, 21, 26] that obstructions to deformations of X are contained in the kernel of

$$\boldsymbol{i}\colon H^2(X,T_X) \to \bigoplus_i \operatorname{Hom}\left(F^pH^i(X,\mathbb{C}), \frac{H^{i+1}(X,\mathbb{C})}{F^pH^{i+1}(X,\mathbb{C})}\right), \qquad \boldsymbol{i}_\xi(\omega) = \xi \lrcorner \omega$$

(this fact is known as Kodaira's Principle *ambient cohomology annihilates obstruction*).
However the proofs of [3, 21, 26] are not completely satisfying because the period map plays only a marginal role in them, while the most natural way of proving Kodaira's Principle would be to show that the period map is a "morphism of deformation theories", i.e. that $\mathcal{P}^p$ commutes with every deformation theoretic construction: for instance obstruction theories.
It is well known that the deformations of a compact complex manifold X are governed

by the L_∞-algebra underlying the Kodaira-Spencer algebra $K_X := \oplus_i A_X^{0,i}(T_X)$.

The main results of this paper are:

1. We explicitly describe an L_∞-algebra C^p such that
$$\{\phi\colon \operatorname{Spec}(B) \to \operatorname{Grass}(H^*(X,\mathbb{C})) \mid \phi(\operatorname{Spec}(\mathbb{C})) = F^pH^*(X,\mathbb{C})\} = \operatorname{Def}_{C^p}(B).$$
2. We explicitly describe a linear L_∞-morphism $K_X \to C^p$ inducing $\mathcal{P}^p$ at the level of deformation functors.

These results give us an algebraic description of the period map and imply, by general theory of L_∞-algebras, that $\mathcal{P}^p$ is a morphism of deformation theories.

The paper is divided in three parts: roughly speaking, in the first we make a functorial construction of an L_∞ structure on the mapping cone of a morphism of differential graded Lie algebras and we introduce the notion of Cartan homotopy. A slightly expanded version of this part is available as [5].
In the second part we describe an L_∞-algebra governing deformations of subcomplexes and then giving a local description of Grassmannians.
Finally in the third part we exhibit an L_∞-morphism inducing the universal period map of a compact Kähler manifolds.

Acknowledgment. Our thanks to Jim Stasheff for precious comments on the first version of [5] and to Barbara Fantechi for useful discussions about dg-Grassmannians.

Keywords and general notation.
We assume that the reader is familiar with the notion and main properties of differential graded Lie algebras and L_∞-algebras (we refer to [6, 9, 14, 17, 18, 22] as introduction of such structures); however the basic definitions are recalled in this paper in order to fix notation and terminology.
For the whole paper, $\mathbb{K}$ is a field of characteristic 0; every vector space is intended over $\mathbb{K}$.
Art is the category of local Artinian $\mathbb{K}$-algebras with residue field $\mathbb{K}$. For $A \in$ **Art** we denote by $\mathfrak{m}_A$ the maximal ideal of A.

Part 1. L_∞ structures on mapping cones

Let $\chi\colon L \to M$ be a morphism of differential graded Lie algebras over a field $\mathbb{K}$ of characteristic 0. In the paper [23] one of the authors has introduced, having in mind the example of embedded deformations, the notion of Maurer-Cartan equation and gauge action for the triple (L, M, χ); these notions reduce to the standard Maurer-Cartan equation and gauge action of L when $M = 0$. More precisely there are defined two functors of Artin rings $\operatorname{MC}_\chi, \operatorname{Def}_\chi\colon$ **Art** $\to$ **Set**, where **Art** is the category of local Artinian $\mathbb{K}$-algebras with residue field $\mathbb{K}$, in the following way:

$$\operatorname{MC}_\chi(A) = \left\{(x, e^a) \in (L^1 \otimes \mathfrak{m}_A) \times \exp(M^0 \otimes \mathfrak{m}_A) \mid dx + \frac{1}{2}[x,x] = 0,\ e^a * \chi(x) = 0\right\},$$

where for $a \in M^0 \otimes \mathfrak{m}_A$ we denote by $\operatorname{ad}_a\colon M\otimes\mathfrak{m}_A \to M\otimes\mathfrak{m}_A$ the operator $\operatorname{ad}_a(y) = [a,y]$ and

$$e^a * y = y + \sum_{n=0}^{+\infty} \frac{\operatorname{ad}_a^n}{(n+1)!}([a,y] - da) = y + \frac{e^{\operatorname{ad}_a} - 1}{\operatorname{ad}_a}([a,y] - da), \qquad y \in M^1 \otimes \mathfrak{m}_A.$$

Then one defines

$$\mathrm{Def}_\chi(A)=\frac{\mathrm{MC}_\chi(A)}{\text{gauge equivalence}},$$

where two solutions of the Maurer-Cartan equation are gauge equivalent if they belong to the same orbit of the gauge action

$$(\exp(L^0\otimes\mathfrak{m}_A)\times\exp(dM^{-1}\otimes\mathfrak{m}_A))\times\mathrm{MC}_\chi(A)\xrightarrow{*}\mathrm{MC}_\chi(A)$$

given by the formula

$$(e^l,e^{dm})*(x,e^a)=(e^l*x,e^{dm}e^ae^{-\chi(l)})=(e^l*x,e^{dm\bullet a\bullet(-\chi(l))}).$$

The $\bullet$ in the rightmost term in the above formula is the Baker-Campbell-Hausdorff multiplication; namely $e^xe^y=e^{x\bullet y}$.
Several examples in [23] illustrate the utility of this construction in deformation theory. In the same paper it is also proved that Def_χ is the truncation of an extended deformation functor [20, Def. 2.1] F such that $T^iF=H^i(C_\chi)$, where C_χ is the differential graded vector space $C_\chi=(\oplus_i C^i_\chi,\delta)$,

$$C^i_\chi=L^i\oplus M^{i-1},\qquad \delta(l,m)=(dl,\chi(l)-dm),\qquad l\in L, m\in M.$$

By a general result [20, Thm. 7.1] there exists an L_∞ structure on C_χ, defined up to homotopy, whose associated deformation functor is isomorphic to Def_χ.
The main result of this part is to describe explicitly a canonical (and hence functorial) L_∞ structure on C_χ with the above property; this is done in an elementary way, without using the theory of extended deformation functors, so that the result of this part can be used to reprove the main results of [23] without using [20].

1. CONVENTIONS ON GRADED VECTOR SPACES

In this paper we will work with $\mathbb{Z}$-graded vector spaces; we write a graded vector space as $V=\oplus_{n\in\mathbb{Z}}V^n$, and call V^n the degree n component of V; an element v of V^n is called a degree n homogeneous element of V.

We adopt the convention according to which degrees are 'shifted on the left'. By this we mean that, for every integer n, $V[n]\simeq\mathbb{K}[n]\otimes V$ where $\mathbb{K}[n]$ denotes the graded vector space consisting in the field $\mathbb{K}$ concentrated in degree $-n$. Note that, with this convention the canonical isomorphism $V\otimes\mathbb{K}[n]\simeq V[n]$ is $v\otimes 1_{[n]}\mapsto(-1)^{n\deg(v)}v_{[n]}$ and we have the following isomorphism, usually called *decalage*

$$\begin{aligned}V_1[1]\otimes\cdots\otimes V_n[1]&\xrightarrow{\sim}(V_1\otimes\cdots\otimes V_n)[n]\\ v_{1[1]}\otimes\cdots\otimes v_{n[1]}&\mapsto(-1)^{\sum_{i=1}^n(n-i)\cdot\deg v_i}(v_1\otimes\cdots\otimes v_n)_{[n]}.\end{aligned}$$

Denote by $\bigotimes^n V$, $\bigodot^n V$ and $\bigwedge^n V$ the n-th tensor, symmetric and exterior powers of V respectively. As with ordinary vector spaces, one can identify $\bigodot^n V$ and $\bigwedge^n V$ with suitable subspaces of $\bigotimes^n V$, called the subspace of symmetric and antisymmetric tensors respectively. The decalage induces a canonical isomorphism

$$\bigodot^n(V[1])\xrightarrow{\sim}\left(\bigwedge^n V\right)[n].$$

Remark 1.1. Using the natural isomorphisms

$$\mathrm{Hom}^i(V,W[l])\simeq\mathrm{Hom}^{i+l}(V,W)$$

and the decalage isomorphism, we obtain natural identifications

$$\mathrm{dec}\colon \mathrm{Hom}^i\left(\bigotimes^k V, W\right) \xrightarrow{\sim} \mathrm{Hom}^{i+k-l}\left(\bigotimes^k (V[1]), W[l]\right),$$

where

$$\mathrm{dec}(f)(v_{1[1]}\otimes\cdots\otimes v_{k[1]}) = (-1)^{ki+\sum_{j=1}^k (k-j)\cdot\deg(v_j)} f(v_1\otimes\cdots\otimes v_k)_{[l]}.$$

By the above considerations

$$\mathrm{dec}\colon \mathrm{Hom}^i\left(\bigwedge^k V, V\right) \xrightarrow{\sim} \mathrm{Hom}^{i+k-1}\left(\bigodot^k (V[1]), V[1]\right).$$

2. Differential graded Lie algebras and L_∞-algebras

A differential graded Lie algebra (DGLA for short) is a Lie algebra in the category of graded vector spaces, endowed with a compatible degree 1 differential. More explicitly, it is the data $(V, d, [\,,\,])$, where V is a graded vector space, the Lie bracket

$$[\,,\,]\colon V\wedge V \to V$$

satisfies the graded Jacobi identity:

$$[v_1,[v_2,v_3]] = [[v_1,v_2],v_3] + (-1)^{\deg(v_1)\deg(v_2)}[v_2,[v_1,v_3]],$$

and $d\colon V\to V$ is a degree 1 differential, i.e.,

$$d^2 = 0, \qquad d[v_1,v_2] = [dv_1,v_2] + (-1)^{\deg(v_1)}[v_1,dv_2].$$

Via the decalage isomorphisms one can look at the Lie bracket of a DGLA V as a morphism

$$q_2\in\mathrm{Hom}^1(V[1]\odot V[1], V[1]), \qquad q_2(v_{[1]}\odot w_{[1]}) = (-1)^{\deg(v)}[v,w]_{[1]},$$

Similarly, the suspended differential $q_1 = d_{[1]} = \mathrm{id}_{\mathbb{K}[1]}\otimes d$ is a morphism of degree 1

$$q_1\colon V[1]\to V[1], \qquad q_1(v_{[1]}) = -(dv)_{[1]}.$$

Up to the canonical bijective linear map $V\to V[1]$, $v\mapsto v_{[1]}$, the suspended differential q_1 and the bilinear operation q_2 are written simply as

$$q_1(v) = -dv; \qquad q_2(v\odot w) = (-1)^{\deg_V(v)}[v,w].$$

Define morphisms $q_k\in\mathrm{Hom}^1(\odot^k(V[1]), V[1])$ by setting $q_k\equiv 0$, for $k\geq 3$. The map

$$\sum_{n\geq 1} q_n\colon \bigoplus_{n\geq 1}\bigodot^n V[1]\to V[1]$$

extends to a coderivation of degree 1

$$Q\colon \bigoplus_{n\geq 1}\bigodot^n V[1] \to \left(\bigoplus_{n\geq 1}\bigodot^n V[1]\right)$$

on the reduced symmetric coalgebra cogenerated by $V[1]$, by the formula

2.1.

$$Q(v_1\odot\cdots\odot v_n) = \sum_{k=1}^n \sum_{\sigma\in S(k,n-k)} \varepsilon(\sigma) q_k(v_{\sigma(1)}\odot\cdots\odot v_{\sigma(k)})\odot v_{\sigma(k+1)}\odot\cdots\odot v_{\sigma(n)},$$

where $S(k,n-k)$ is the set of unshuffles and $\varepsilon(\sigma)=\pm 1$ is the *Koszul sign*, determined by the relation in $\bigodot^n V[1]$

$$v_{\sigma(1)}\odot\cdots\odot v_{\sigma(n)}=\varepsilon(\sigma)v_1\odot\cdots\odot v_n.$$

The axioms of differential graded Lie algebra are then equivalent to Q being a codifferential, i.e., $Q^2=0$. This description of differential graded Lie algebras in terms of the codifferential Q is called the Quillen construction [25]. By dropping the requirement that $q_k\equiv 0$ for $k\geq 3$ one obtains the notion of L_∞-algebra (or strong homotopy Lie algebra), see e.g. [17, 18, 14]; namely, an L_∞ structure on a graded vector space V is a sequence of linear maps of degree 1

$$q_k\colon \bigodot^k V[1]\to V[1],\qquad k\geq 1,$$

such that the induced coderivation Q on the reduced symmetric coalgebra cogenerated by $V[1]$, given by the Formula 2.1 is a codifferential, i.e. i.e., $Q^2=0$. This condition in particular implies $q_1^2=0$, i.e., an L_∞-algebra is in particular a differential complex. Note that, by the above discussion, every DGLA can be naturally seen as an L_∞-algebra; namely, a DGLA is an L_∞-algebra with vanishing higher multiplications q_k, $k\geq 3$. Note that, via the decalage isomorphisms of Remark 1.1, every component q_k of an L_∞ structure on V can be seen as morphism

$$\mu_k\in \operatorname{Hom}^{2-k}(\bigwedge^k V,V).$$

A morphism f_∞ between two L_∞-algebras $(V,q_1,q_2,q_3,\dots)$ and $(W,p_1,p_2,p_3,\dots)$ is a sequence of linear maps of degree 0

$$f_n\colon \bigodot^n V[1]\to W[1],\qquad n\geq 1,$$

such that the morphism of coalgebras

$$F\colon \bigoplus_{n\geq 1}\bigodot^n V[1]\to \bigoplus_{n\geq 1}\bigodot^n W[1]$$

induced by $\sum_n f_n\colon \bigoplus_{n\geq 1}\bigodot^n V[1]\to W[1]$ commutes with the codifferentials induced by the two L_∞ structures on V and W [6, 14, 17, 18, 22]. An L_∞-morphism f_∞ is called *linear* (sometimes *strict*) if $f_n=0$ for every $n\geq 2$. We note that a linear map $f_1\colon V[1]\to W[1]$ is a linear L_∞-morphism if and only if

$$p_n(f_1(v_1)\odot\cdots\odot f_1(v_n))=f_1(q_n(v_1\odot\cdots\odot v_n)),\qquad \forall\, n\geq 1,\ v_1,\dots,v_n\in V[1].$$

For instance, morphisms between DGLAs are linear morphisms between the corresponding L_∞-algebras.

If f_∞ is an L_∞-morphism between $(V,q_1,q_2,q_3,\dots)$ and $(W,p_1,p_2,p_3,\dots)$, then its linear part

$$f_1\colon V[1]\to W[1]$$

satisfies the equation $f_1\circ q_1=p_1\circ f_1$, i.e., f_1 is a map of differential complexes $(V[1],q_1)\to(W[1],p_1)$. An L_∞-morphism f_∞ is called a quasiisomorphism of L_∞-algebras if its linear part f_1 is a quasiisomorphism of differential complexes. A major result in the theory of L_∞-algebra is the following *homotopical transfer of structure* theorem (see [6, 16] for a proof).

Theorem 2.2. *Let* $(V,q_1,q_2,q_3,\dots)$ *be an* L_∞*-algebra and* (C,δ) *be a differential complex. If there exist two morphisms of differential complexes*

$$\imath\colon (C[1],\delta_{[1]})\to (V[1],q_1)\qquad \text{and}\qquad \pi\colon (V[1],q_1)\to (C[1],\delta_{[1]})$$

which are homotopy inverses, then there exist an L_∞*-algebra structure* $(C,\langle\,\rangle_1,\langle\,\rangle_2,\dots)$ *on* C *extending its differential complex structure, and making* $(V,q_1,q_2,\dots)$ *and* $(C,\langle\,\rangle_1,\langle\,\rangle_2,\dots)$ *be quasiisomorphic* L_∞*-algebras via an* L_∞*-quasiisomorphism* $\imath_\infty$ *extending* $\imath$.

In case $\pi\imath=\mathrm{Id}_{C[1]}$, explicit formulas for such a transfer are described in [6, 16] and [32] in terms of summation over rooted trees [15, Definition 6]

$$\langle\,\rangle_n=\sum_{\Gamma\in T_n}\varepsilon_\Gamma\frac{Z_\Gamma(\imath,\pi,K,q_i)}{|\operatorname{Aut}\Gamma|},$$

where $K\in\operatorname{Hom}^{-1}(V[1],V[1])$ is an homotopy between $\imath\pi$ and $\mathrm{Id}_{V[1]}$, T_n is the set of rooted trees with n tails, $\varepsilon_\Gamma=\pm1$ is a sign depending on the combinatorics of the tree Γ and $\operatorname{Aut}\Gamma$ is the group of automorphisms of Γ. Each tail edge of a tree is decorated by the operator $\imath$, each internal edge is decorated by the suspended operator K and the root edge is decorated by the suspended operator π; every internal vertex v carries the operation q_r, where r is the number of edges having v as endpoint. Then Z_Γ is the evaluation of such a decorated graph according to the usual operadic rules; see [6, Thm. 2.3.1] for an explicit recursive formula.

3. The suspended mapping cone of $\chi\colon L\to M$.

The suspended mapping cone of the DGLA morphism $\chi\colon L\to M$ is the graded vector space

$$C_\chi=\operatorname{Cone}(\chi)[-1],$$

where $\operatorname{Cone}(\chi)=L[1]\oplus M$ is the mapping cone of χ. More explicitly,

$$C_\chi=\bigoplus_i C_\chi^i,\qquad C_\chi^i=L^i\oplus M^{i-1}.$$

The suspended mapping cone has a natural differential $\delta\in\operatorname{Hom}^1(C_\chi,C_\chi)$ given by

$$\delta(l,m)=(dl,\chi(l)-dm),\qquad l\in L, m\in M.$$

Denote by $\langle\,\rangle_1\in\operatorname{Hom}^1(C_\chi[1],C_\chi[1])$ the suspended differential, namely

$$\langle(l,m)\rangle_1=(-dl,-\chi(l)+dm),\qquad l\in L, m\in M.$$

Remark 3.1. If χ is injective, then the projection on the second factor induces a quasiisomorphism of differential complexes $\pi_2\colon C_\chi\to(M/\operatorname{Im}(\chi))[-1]$. In particular, it induces isomorphisms $H^i(C_\chi)\simeq H^{i-1}(\operatorname{Coker}(\chi))$, for every i.

Setting $M[t,dt]=\mathbb{K}[t,dt]\otimes M$, then

$$H_\chi=\{(l,m(t,dt))\in L\times M[t,dt]\mid m(0,0)=0,\ m(1,0)=\chi(l)\}$$

is a differential graded Lie algebra. The differential on H_χ is $(l,m(t,dt))\mapsto(dl,dm(t,dt))$; since the differential on H_χ has degree 1, the suspended differential $q_1\colon H_\chi[1]\to H_\chi[1]$ is the opposite differential:

$$q_1(l,m(t,dt))=-(dl,dm(t,dt)).$$

The integral operator $\int_a^b\colon \mathbb{K}[t,dt]\to\mathbb{K}$ extends naturally to a linear map of degree -1

$$\int_a^b\colon M[t,dt]\to M,\qquad \int_a^b(\sum_i t^i m_i + t^i dt\cdot n_i) = \sum_i\left(\int_a^b t^i dt\right) n_i.$$

Lemma 3.2. *The complexes* $C_\chi[1]$ *and* $H_\chi[1]$ *are homotopic; more precisely, if one denotes by*

$$\imath\in \operatorname{Hom}^0(C_\chi[1],H_\chi[1]),\qquad \pi\in\operatorname{Hom}^0(H_\chi[1],C_\chi[1]),\qquad K\in\operatorname{Hom}^{-1}(H_\chi[1],H_\chi[1])$$

the linear maps defined as

$$\imath(l,m)=(l,t\chi(l)+dt\cdot m),\qquad \pi(l,m(t,dt))=\left(l,\int_0^1 m(t,dt)\right)$$

$$K(l,m)=\left(0,t\int_0^1 m-\int_0^t m\right),$$

then ι *and* π *are morphisms of complexes and*

$$\pi\imath=\operatorname{Id}_{C_\chi[1]},\qquad \operatorname{Id}_{H_\chi[1]}-\imath\pi = Kq_1+q_1K.$$

Proof. Straightforward. □

4. THE L_∞ STRUCTURE ON C_χ

By Quillen construction [25], the differential graded Lie algebra H_χ carries an L_∞ structure

$$q_k\colon \overset{k}{\bigodot}(H_\chi[1])\to H_\chi[1],$$

where $q_k=0$ for every $k\geq 3$,

$$q_1(l,m(t,dt))=(-dl,-dm(t,dt))$$

and

$$q_2((l_1,m_1(t,dt))\odot(l_2,m_2(t,dt)))=(-1)^{\deg_{H_\chi}(l_1,m_1(t,dt))}([l_1,l_2],[m_1(t,dt),m_2(t,dt)]).$$

Results of Section 3 tell us that we can apply the homotopy transfer of structure theorem, to induce on C_χ an L_∞-algebra structure making C_χ and H_χ be quasiisomorphic L_∞-algebras. Moreover, the linear maps of degree 1

$$\langle\ \rangle_n\colon \overset{n}{\bigodot} C_\chi[1]\to C_\chi[1],\qquad n\geq 1,$$

defining the induced L_∞-algebra structure on C_χ are explicitly described in terms of summation over rooted trees. In our case, the properties

$$\pi q_1 K = K q_1\imath = 0,$$

$$q_2(\operatorname{Im}K\otimes\operatorname{Im}K)\subseteq\ker\pi\cap\ker K,\qquad q_k=0\ \ \forall\, k\geq 3,$$

imply that, fixing the number of tails, there exists at most one isomorphism class of trees giving a nontrivial contribution.

- One tail: the only tree is

giving by operadic evaluation the formula

$$\langle(l,m)\rangle_1=\pi q_1\imath(l,m)=(-dl,-\chi(l)+dm).$$

- Two tails:



Again by operadic evaluation, this graph gives

$$\langle \gamma_1 \odot \gamma_2 \rangle_2 = \pi q_2(\imath(\gamma_1) \odot \imath(\gamma_2)).$$

- n tails:



This diagram gives, for every $n \geq 2$ the formula

$$\langle \gamma_1 \odot \cdots \odot \gamma_n \rangle_n =$$
$$= \frac{(-1)^{n-2}}{2} \sum_{\sigma \in S_n} \varepsilon(\sigma) \pi q_2(\imath(\gamma_{\sigma(1)}) \odot K q_2(\imath(\gamma_{\sigma(2)}) \odot \cdots \odot K q_2(\imath(\gamma_{\sigma(n-1)}) \odot \imath(\gamma_{\sigma(n)})) \cdots)).$$

The factor 1/2 in the above formula accounts for the cardinality of the automorphism group of the graph involved.

Remark 4.1. The above construction of the L_∞ structure on C_χ commutes with tensor products of differential graded commutative algebras. This means that if R is a DGCA, then the L_∞-algebra structure on the suspended mapping cone of $\chi \otimes \mathrm{id}_{\mathrm{R}} \colon L \otimes R \to M \otimes R$ is naturally isomorphic to the L_∞-algebra $C_\chi \otimes R$.

A more refined description involving the original brackets in the differential graded Lie algebras L and M is obtained decomposing the symmetric powers of $C_\chi[1]$ into types:

$$\bigodot^n (C_\chi[1]) = \bigodot^n \mathrm{Cone}(\chi) = \bigoplus_{\lambda+\mu=n} \left(\bigodot^\mu M \right) \otimes \left(\bigodot^\lambda L[1] \right).$$

The operation $\langle\,\rangle_2$ decomposes into

$$l_1 \otimes l_2 \mapsto (-1)^{\deg_L(l_1)}[l_1, l_2] \in L; \qquad m_1 \otimes m_2 \mapsto 0;$$

$$m \otimes l \mapsto \frac{(-1)^{\deg_M(m)+1}}{2}[m, \chi(l)] \in M.$$

For later use, we point out that, via decalage isomorphisms, the maps $\langle - \rangle_1$ and $\langle - \rangle_2$ corresponds to

$$\mu_1 \in \mathrm{Hom}^1(C_\chi, C_\chi), \qquad \mu_2 \in \mathrm{Hom}^0(\bigwedge^2 C_\chi, C_\chi),$$

$$\mu_1(l, m) = (dl, \chi(l) - dm),$$

$$\mu_2((l_1,m_1)\wedge(l_2,m_2))=\left([l_1,l_2],\frac{1}{2}[m_1,\chi(l_2)]+\frac{(-1)^{\deg_L(l_1)}}{2}[\chi(l_1),m_2]\right).$$

For every $n\geq 2$ it is easy to see that $\langle\gamma_1\odot\cdots\odot\gamma_{n+1}\rangle_{n+1}$ can be nonzero only if the multivector $\gamma_1\odot\cdots\odot\gamma_{n+1}$ belongs to $\bigodot^n M\otimes L[1]$. For $n\geq 2$, $m_1,\ldots,m_n\in M$ and $l\in L[1]$ the formula for $\langle-\rangle_{n+1}$ described above becomes

$$\begin{aligned}&\langle m_1\odot\cdots\odot m_n\odot l\rangle_{n+1}=\\&=(-1)^{n-1}\sum_{\sigma\in S_n}\varepsilon(\sigma)\pi q_2((dt)m_{\sigma(1)}\odot Kq_2((dt)m_{\sigma(2)}\odot\cdots\odot Kq_2((dt)m_{\sigma(n)}\odot t\chi(l))\cdots)).\end{aligned}$$

Define recursively a sequence of polynomials $\phi_i(t)\in\mathbb{Q}[t]\subseteq\mathbb{K}[t]$ and rational numbers I_n by the rule

$$\phi_1(t)=t,\qquad I_n=\int_0^1\phi_n(t)dt,\qquad \phi_{n+1}(t)=\int_0^t\phi_n(s)ds-tI_n.$$

By the definition of the homotopy operator K we have, for every $m\in M$

$$K((\phi_n(t)dt)m)=-\phi_{n+1}(t)m.$$

Therefore, for every $m_1,m_2\in M$ we have

$$Kq_2((dt\cdot m_1)\odot\phi_n(t)m_2)=(-1)^{\deg_M(m_1)}\phi_{n+1}(t)[m_1,m_2].$$

Therefore, we find:

$$\begin{aligned}&\langle m_1\odot\cdots\odot m_n\odot l\,\rangle_{n+1}=\\&=(-1)^{n-1}\sum_{\sigma\in S_n}\varepsilon(\sigma)\pi q_2((dt)m_{\sigma(1)}\odot Kq_2((dt)m_{\sigma(2)}\odot\cdots\odot Kq_2((dt)m_{\sigma(n)}\odot t\chi(l))\cdots))\\&=(-1)^{n-1+\deg_M(m_{\sigma(n)})}\sum_{\sigma\in S_n}\varepsilon(\sigma)\pi q_2((dt)m_{\sigma(1)}\odot Kq_2((dt)m_{\sigma(2)}\odot\cdots\odot\phi_2(t)[m_{\sigma(n)},\chi(l)]\cdots))\\&=(-1)^{n-1+\sum_{i=2}^n\deg_M(m_{\sigma(i)})}\sum_{\sigma\in S_n}\varepsilon(\sigma)\pi q_2((dt)m_{\sigma(1)}\odot\phi_n(t)[m_{\sigma(2)},\cdots,[m_{\sigma(n)},\chi(l)]\cdots])\\&=(-1)^{n+\sum_{i=1}^n\deg_M(m_i)}I_n\sum_{\sigma\in S_n}\varepsilon(\sigma)[m_{\sigma(1)},[m_{\sigma(2)},\cdots,[m_{\sigma(n)},\chi(l)]\cdots]]\in M\end{aligned}$$

Theorem 4.2. *For every $n\geq 2$ we have*

$$\begin{aligned}&\langle m_1\odot\cdots\odot m_n\odot l\,\rangle_{n+1}=\\&\qquad=-(-1)^{\sum_{i=1}^n\deg_M(m_i)}\frac{B_n}{n!}\sum_{\sigma\in S_n}\varepsilon(\sigma)[m_{\sigma(1)},[m_{\sigma(2)},\cdots,[m_{\sigma(n)},\chi(l)]\cdots]],\end{aligned}$$

where the B_n are the Bernoulli numbers, defined by the series expansion identity

$$\sum_{n=0}^{\infty}B_n\frac{x^n}{n!}=\frac{x}{e^x-1}=1-\frac{x}{2}+\frac{x^2}{12}-\frac{x^4}{720}+\cdots$$

Proof. Since $B_{2k+1}=0$ for every $k>0$, it is sufficient to prove that $B_n=-n!I_n$ for every $n\geq 1$. Consider the polynomials $\psi_0(t)=1$ and $\psi_n(t)=n!(\phi_n(t)-I_n)$ for $n\geq 1$. Then, for any $n\geq 1$,

$$\frac{d}{dt}\psi_n(t)=n\cdot\psi_{n-1}(t),\qquad \int_0^1\psi_n(t)dt=0.$$

Therefore the $\psi_n(t)$ satisfy the recursive relations (see e.g. [28]) of the Bernoulli polynomials $B_n(t)$, defined by the series expansion identity

$$\sum_{n=0}^{\infty} B_n(t)\frac{x^n}{n!} = \frac{xe^{tx}}{e^x-1}.$$

In particular $B_n = B_n(0) = \psi_n(0) = -n!I_n$ for every $n \geq 1$. □

Remark 4.3. Recently, the relevance of Bernoulli numbers in deformation theory has been also remarked by Ziv Ran in [27]. In particular, Ran's "JacoBer" complex seems to be closely related to the coderivation Q defining the L_∞ structure on C_χ.

5. The functors MC_χ and Def_χ revisited

Having introduced an L_∞ structure on C_χ in Section 4, we have a corresponding Maurer-Cartan functor [6, 14] $\mathrm{MC}_{C_\chi}\colon \mathbf{Art}\to\mathbf{Set}$, defined as

$$\mathrm{MC}_{C_\chi}(A) = \left\{ \gamma \in C_\chi[1]^0\otimes\mathfrak{m}_A \;\middle|\; \sum_{n\geq 1}\frac{\langle\gamma^{\odot n}\rangle_n}{n!} = 0 \right\}, \qquad A\in\mathbf{Art}.$$

Writing $\gamma = (l,m)$, with $l \in L^1\otimes\mathfrak{m}_A$ and $m\in M^0\otimes\mathfrak{m}_A$, the Maurer-Cartan equation becomes

$$\begin{aligned}
0 &= \sum_{n=1}^{\infty}\frac{\langle (l,m)^{\odot n}\rangle_n}{n!}\\
&= \langle (l,m)\rangle_1 + \frac{1}{2}\langle l^{\odot 2}\rangle_2 + \langle m\otimes l\rangle_2 + \frac{1}{2}\langle m^{\odot 2}\rangle_2 + \sum_{n\geq 2}\frac{n+1}{(n+1)!}\langle m^{\odot n}\otimes l\rangle_{n+1}\\
&= \left(-dl - \frac{1}{2}[l,l], -\chi(l) + dm - \frac{1}{2}[m,\chi(l)] + \sum_{n\geq 2}\frac{1}{n!}\langle m^{\odot n}\otimes l\rangle_{n+1}\right) \in (L^2\oplus M^1)\otimes\mathfrak{m}_A.
\end{aligned}$$

According to Theorem 4.2, since $\deg_M(m) = \deg_{C_\chi[1]}(m) = 0$, we have

$$\langle m^{\odot n}\otimes l\rangle_{n+1} = -\frac{B_n}{n!}\sum_{\sigma\in S_n}[m,[m,\cdots,[m,\chi(l)]\cdots]] = -B_n\,\mathrm{ad}_m^n(\chi(l)).$$

The Maurer-Cartan equation on C_χ is therefore equivalent to

$$\begin{cases} dl + \dfrac{1}{2}[l,l] = 0 \\[2ex] \chi(l) - dm + \dfrac{1}{2}[m,\chi(l)] + \displaystyle\sum_{n=2}^{\infty}\frac{B_n}{n!}\,\mathrm{ad}_m^n(\chi(l)) = 0. \end{cases}$$

Since $B_0 = 1$ and $B_1 = -\dfrac{1}{2}$, we can write the second equation as

$$[m,\chi(l)] - dm + \sum_{n=0}^{\infty}\frac{B_n}{n!}\,\mathrm{ad}_m^n(\chi(l)) = [m,\chi(l)] - dm + \frac{\mathrm{ad}_m}{e^{\mathrm{ad}_m}-1}(\chi(l)) = 0.$$

Applying the invertible operator $\dfrac{e^{\mathrm{ad}_m}-1}{\mathrm{ad}_m}$ we get

$$0=\chi(l)+\frac{e^{\mathrm{ad}_m}-1}{\mathrm{ad}_m}([m,\chi(l)]-dm)=e^m*\chi(l).$$

Therefore, the Maurer-Cartan equation for the L_∞-algebra structure on C_χ is equivalent to

$$\begin{cases} dl+\dfrac{1}{2}[l,l]=0\\ e^m*\chi(l)=0\end{cases}$$

and the Maurer-Cartan functor MC_χ described in the introduction is precisely the Maurer-Cartan functor corresponding to the L_∞ structure on $C\chi$.

Recall that the deformation functor associated to an L_∞-algebra $\mathfrak{g}$ is $\mathrm{Def}_{\mathfrak{g}}=\mathrm{MC}_{\mathfrak{g}}/\sim$, where $\sim$ denotes homotopy equivalence of solutions of the Maurer-Cartan equation: two elements γ_0 and γ_1 of $\mathrm{MC}_{\mathfrak{g}}(A)$ are called homotopy equivalent if there exists an element $\gamma(t,dt)\in\mathrm{MC}_{\mathfrak{g}[t,dt]}(A)$ with $\gamma(0)=\gamma_0$ and $\gamma(1)=\gamma_1$.

We have already identified the functor MC_{C_χ} with the functor MC_χ. Now we want to show that, under this identification, the homotopy equivalence on MC_{C_χ} is the same thing of gauge equivalence on MC_χ described in the introduction, so that

$$\mathrm{Def}_\chi\simeq\mathrm{Def}_{C_\chi}.$$

We will need the following lemma (see the Appendix A for a proof).

Lemma 5.1. *Let* $\mathfrak{g}$ *be a differential graded Lie algebra and let* $A\in\mathbf{Art}$. *Then, for any* x *in* $\mathrm{MC}_{\mathfrak{g}}(A)$ *and any* $g(t)\in\mathfrak{g}^0[t]\otimes\mathfrak{m}_A$, *with* $g(0)=0$, *the element* $e^{g(t)}*x$ *is an element of* $\mathrm{MC}_{\mathfrak{g}[t,dt]}(A)$. *Moreover all the elements of* $\mathrm{MC}_{\mathfrak{g}[t,dt]}(A)$ *are obtained in this way.*

We first show that homotopy implies gauge. Let (l_0,m_0) and (l_1,m_1) be homotopy equivalent elements of $\mathrm{MC}_{C_\chi}(A)$. Then there exists an element $(\tilde{l},\tilde{m})$ of $\mathrm{MC}_{C_\chi[s,ds]}(A)$ with $(\tilde{l}(0),\tilde{m}(0))=(l_0,m_0)$ and $(\tilde{l}(1),\tilde{m}(1))=(l_1,m_1)$. According to Remark 4.1, the Maurer-Cartan equation for $(\tilde{l},\tilde{m})$ is

$$\begin{cases} d\tilde{l}+\dfrac{1}{2}[\tilde{l},\tilde{l}]=0\\ e^{\tilde{m}}*\chi(\tilde{l})=0\end{cases}$$

The first of the two equations above tells us that $\tilde{l}$ is a solution of the Maurer-Cartan equation for $L[s,ds]$. So, by Lemma 5.1, there exists a degree zero element $\lambda(s)$ in $L[s]\otimes\mathfrak{m}_A$ with $\lambda(0)=0$ such that $\tilde{l}=e^\lambda*l_0$. Evaluating at $s=1$ we find $l_1=e^{\lambda_1}*l_0$. As a consequence of $\tilde{l}=e^\lambda*l_0$, we also have $\chi(\tilde{l})=e^{\chi(\lambda)}*\chi(l_0)$. Set $\tilde{\mu}=\tilde{m}\bullet\chi(\lambda)\bullet m_0$, so that $\tilde{m}=\tilde{\mu}\bullet m_0\bullet(-\chi(\lambda))$ and the second Maurer-Cartan equation is reduced to $e^{\tilde{\mu}}*(e^{m_0}*\chi(l_0))=0$, i.e., to $e^{\tilde{\mu}}*0=0$, where we have used the fact that (l_0,m_0) is a solution of the Maurer-Cartan equation in C_χ. This last equation is equivalent to the equation $d\tilde{\mu}=0$ in $(C_\chi[s,ds])^0\otimes\mathfrak{m}_A$. If we write $\tilde{\mu}(s,ds)=\mu^0(s)+ds\,\mu^{-1}(s)$, then the equation $d\tilde{\mu}=0$ becomes

$$\begin{cases}\dot{\mu}^0-d_M\mu^{-1}=0\\ d_M\mu^0=0,\end{cases}$$

where d_M is the differential in the DGLA M. The solution is, for any fixed μ^{-1},

$$\mu^0(s)=\int_0^s d\sigma\,d_M\mu^{-1}(\sigma)=-d_M\int_0^s d\sigma\,\mu^{-1}(\sigma)$$

Set $\nu = -\int_0^1 ds\, \mu^{-1}(s)$. Then $m_1 = \tilde{m}(1) = (d_M\nu) \bullet m_0 \bullet (-\chi(\lambda_1))$. Summing up, if (l_0, m_0) and (m_1, l_1) are homotopy equivalent, then there exists $(d\nu, \lambda_1) \in (dM^{-1} \otimes \mathfrak{m}_A) \times (L^0 \otimes \mathfrak{m}_A)$ such that

$$\begin{cases} l_1 = e^{\lambda_1} * l_0 \\ m_1 = d\nu \bullet m_0 \bullet (-\chi(\lambda_1)), \end{cases}$$

i.e., (l_0, m_0) and (m_1, l_1) are gauge equivalent.

We now show that gauge implies homotopy. Assume (l_0, m_0) and (m_1, l_1) are gauge equivalent. Then then there exist $(d\nu, \lambda_1) \in (dM^{-1} \otimes \mathfrak{m}) \times (L^0 \otimes \mathfrak{m})$ such that

$$\begin{cases} l_1 = e^{\lambda_1} * l_0 \\ m_1 = d\nu \bullet m_0 \bullet (-\chi(\lambda_1)). \end{cases}$$

Set $\tilde{l}(s, ds) = e^{s\lambda_1} * l_0$. By Lemma 5.1, $\tilde{l}$ satisfies the equation $d\tilde{l} + \frac{1}{2}[\tilde{l}, \tilde{l}] = 0$. Set $\tilde{m} = (d(s\nu)) \bullet m_0 \bullet (-\chi(s\lambda_1))$. Reasoning as above, we find

$$e^{\tilde{m}} * \chi(\tilde{l}) = e^{d(s\nu)} * 0 = 0.$$

Therefore, $(\tilde{l}, \tilde{m})$ is a solution of the Maurer-Cartan equation in $C_\chi[s, ds]$. Moreover $\tilde{l}(0) = l_0$, $\tilde{l}(1) = l_1$, $\tilde{m}(0) = m_0$ and $\tilde{m}(1) = d\nu \bullet m_0 \bullet (-\chi(\lambda_1)) = m_1$, i.e. (l_0, m_0) and (m_1, l_1) are homotopy equivalent.
Summing up, we have

$$\mathrm{Def}_{C_\chi} = \frac{\mathrm{MC}_{C_\chi}}{\text{homotopy}} \simeq \frac{\mathrm{MC}_\chi}{\text{gauge}} = \mathrm{Def}_\chi\,.$$

6. Functoriality

In the above section we have shown how to a morphism of differential graded Lie algebras $\chi\colon L \to M$ is associated a canonical L_∞ structure on C_χ. We will now discuss the functorial aspects of this construction. Denote by $\mathbf{L}_\infty$ the category of L_∞-algebras and by $\mathbf{M}$ the category of morphisms of differential graded Lie algebras; objects in $\mathbf{M}$ are DGLA morphisms $\chi\colon L \to M$; morphisms in $\mathbf{M}$ are commutative squares

$$\begin{tikzcd} L_1 \arrow[r, "f_L"] \arrow[d, "\chi_1"'] & L_2 \arrow[d, "\chi_2"] \\ M_1 \arrow[r, "f_M"] & M_2 \end{tikzcd}$$

of DGLA morphisms.

It is immediate to observe that the above commutative square induces a linear L_∞-morphism $(f_L, f_M)\colon C_{\chi_1} \to C_{\chi_2}$ and then $C\colon \mathbf{M} \to \mathbf{L}_\infty$ is a functor. Moreover both MC and Def are functors from the category $\mathbf{L}_\infty$ to the category of functors of Artin rings [14, 20, 22]. The functor MC acts on the morphisms of $\mathbf{L}_\infty$ in the following way: let $f_\infty\colon V \to W$ be an L_∞-morphism, then

$$\mathrm{MC}_{f_\infty}\colon\ \mathrm{MC}_V \to \mathrm{MC}_W$$

is the natural transformation given, for $A \in \mathbf{Art}$ and $v \in \mathrm{MC}_V(A) \subseteq V[1]^0 \otimes \mathfrak{m}_A$, by

$$\mathrm{MC}_{f_\infty}(v) = \sum_{n=1}^{\infty} \frac{1}{n!} f_n(v^{\odot n}) \in W[1]^0 \otimes \mathfrak{m}_A.$$

The natural transformation MC_{f_∞} preserves the homotopy equivalence and then induces a natural transformation

$$\mathrm{Def}_{f_\infty}\colon \mathrm{Def}_V \to \mathrm{Def}_W .$$

Recall from Section 2 that an L_∞-morphism f_∞ is called a quasiisomorphism of L_∞-algebras if its linear part f_1 is a quasiisomorphism of differential complexes. Using the fact that every quasiisomorphism of L_∞-algebras induces an isomorphism of the associated deformation functors [14], the next theorem becomes evident.

Theorem 6.1 ([23]). *Consider a commutative diagram of morphisms of differential graded Lie algebras*

$$\begin{tikzcd} L_1 \arrow[r, "f_L"] \arrow[d, "\chi_1"'] & L_2 \arrow[d, "\chi_2"] \\ M_1 \arrow[r, "f_M"] & M_2 \end{tikzcd}$$

and assume that $(f_L, f_M)\colon C_{\chi_1} \to C_{\chi_2}$ *is a quasiisomorphism of complexes (e.g. if both* f_L *and* f_M *are quasiisomorphisms). Then the natural transformation* $\mathrm{Def}_{\chi_1} \to \mathrm{Def}_{\chi_2}$ *is an isomorphism.*

7. Cartan homotopies

In this section we formalize, under the notion of *Cartan homotopy*, a set of standard identities that often arise in algebra and geometry [3, Appendix B].

Definition 7.1. Let $(L, d, [,])$ and $(M, d, [,])$ be two differential graded Lie algebras and denote by δ the standard differential on $\mathrm{Hom}^*(L, M)$. A linear map $i \in \mathrm{Hom}^{-1}(L, M)$ is called a *Cartan homotopy* if for every $a, b \in L$ we have:

$$i([a, b]) = [i(a), \delta i(b)], \qquad [i(a), i(b)] = 0.$$

Notice that, according to the definition of δ, for every $a \in L$ we have

$$\delta i(a) = d(i(a)) + i(da).$$

For later use we point out that $[i(a), [i(b), \delta i(c)]] = [i(a), i([b, c])] = 0$ for every a, b, c. It is moreover easy to verify that δi is a morphism of differential graded Lie algebras and

$$i([a, b]) = (-1)^{\deg(a)}[\delta i(a), i(b)] = \frac{1}{2}[i(a), \delta i(b)] + \frac{(-1)^{\deg(a)}}{2}[\delta i(a), i(b)].$$

Example 7.2. The name Cartan homotopy has a clear origin in differential geometry. Namely, let M be a differential manifold, $\mathcal{X}(M)$ be the Lie algebra of vector fields on M, and $\mathcal{E}nd^*(\Omega^*(M))$ be the Lie algebra of endomorphisms of the de Rham algebra of M. The Lie algebra $\mathcal{X}(M)$ can be seen as a DGLA concentrated in degree zero, and the graded Lie algebra $\mathcal{E}nd^*(\Omega^*(M))$ has a degree one differential given by $[d, -]$, where d is the de Rham differential. Then the contraction

$$i\colon \mathcal{X}(M) \to \mathcal{E}nd^*(\Omega^*(M))[-1]$$

is a Cartan homotopy and its differential is the Lie derivative

$$\delta i = \mathcal{L}\colon \mathcal{X}(M) \to \mathcal{E}nd^*(\Omega^*(M)).$$

In fact, by classical Cartan's homotopy formulas [1, Section 2.4], for any two vector fields X and Y on M, we have

(1) $\mathcal{L}_X = di_X + i_X d = [d, i_X]$;
(2) $i_{[X,Y]} = \mathcal{L}_X i_Y - i_Y \mathcal{L}_X = [\mathcal{L}_X, i_Y] = [i_X, \mathcal{L}_Y]$;
(3) $[i_X, i_Y] = 0$.

Note that the first Cartan formula above actually states that $\delta i = \mathcal{L}$. Indeed $\mathcal{X}(M)$ is concentrated in degree zero and then its differential is trivial.

Example 7.3. The composition of a Cartan homotopy with a morphism of DGLAs is a Cartan homotopy. If $i\colon L \to M[-1]$ is a Cartan homotopy and Ω is a differential graded-commutative algebra, then its natural extension

$$i \otimes \mathrm{Id}\colon L \otimes \Omega \to (M \otimes \Omega)[-1], \qquad a \otimes \omega \mapsto i(a) \otimes \omega,$$

is a Cartan homotopy.

Proposition 7.4. *Let $i\colon L \to M[-1]$ be a Cartan homotopy and $\chi = \delta i\colon L \to M$. Then the linear map*

$$\tilde{i}\colon L \to C_\chi, \qquad \tilde{i}(a) = (a, i(a))$$

is a linear L_∞-morphism.

Proof. By decalage isomorphism, the L_∞ structure on C_χ is given by the higher brackets $\mu_n \in \mathrm{Hom}^{2-n}(\bigwedge^n C_\chi, C_\chi)$, $n \geq 1$, where

$$\mu_1((l,m)) = (dl, \chi(l) - dm),$$

$$\mu_2((l,m) \wedge (h,k)) = \left([l,h], \frac{1}{2}[m, \chi(h)] + \frac{(-1)^{\deg(l)}}{2}[\chi(l), k]\right)$$

and for $n \geq 3$

$$\mu_n((l_1,m_1) \wedge \cdots \wedge (l_n,m_n)) = \frac{B_{n-1}}{(n-1)!} \sum_{\sigma \in S_n} \pm [m_{\sigma(1)}, [\cdots, [m_{\sigma(n-1)}, \chi(l)_{\sigma(n)}] \cdots]]$$

It is straightforward to check that $\tilde{i}$ commutes with every bracket, i.e.

$$\tilde{i}(dx) = \mu_1(\tilde{i}(x)), \qquad \tilde{i}([x,y]) = \mu_2(\tilde{i}(x) \wedge \tilde{i}(y)),$$

and for $n \geq 3$

$$\mu_n(\tilde{i}(x_1) \wedge \cdots \wedge \tilde{i}(x_n)) = 0.$$

Therefore $\tilde{i}$ is a linear L_∞-morphism. □

Appendix A: gauge vs. homotopy

In this Appendix we briefly discuss the relation between homotopy and gauge equivalence for solutions of the Maurer-Cartan equation for a given differential graded Lie algebra L. We also give a proof of Lemma 5.1, which is here presented as Corollary A.1.

Proposition A.1. *Let $(L, d, [\ ,\ ])$ be a differential graded Lie algebra such that:*

(1) *$L = M \oplus C \oplus D$ as graded vector spaces.*
(2) *M is a differential graded subalgebra of L.*
(3) *$d\colon C \to D[1]$ is an isomorphism of graded vector spaces.*

*Then, for every $A \in$ **Art** there exists a bijection*

$$\alpha\colon \mathrm{MC}_M(A) \times (C^0 \otimes \mathfrak{m}_A) \xrightarrow{\sim} \mathrm{MC}_L(A), \qquad (x, c) \mapsto e^c * x.$$

Proof. This is essentially proved in [31, Section 5] using induction on the length of A and the Baker-Campbell-Hausdorff formula.
Here we sketch a simpler proof based on formal theory of deformation functors [30, 29, 4, 19].
The map α is a natural transformation of homogeneous functors, so it is sufficient to show that α is bijective on tangent spaces and injective on obstruction spaces. Recall that the tangent space of MC_L is $Z^1(L)$, while its obstruction space is contained in $H^2(L)$. The functor $A \mapsto C^0 \otimes \mathfrak{m}_A$ is smooth with tangent space C^0 and therefore tangent and obstruction spaces of the functor

$$A \mapsto \mathrm{MC}_M(A) \times (C^0 \otimes \mathfrak{m}_A)$$

are respectively $Z^1(M) \oplus C^0$ and $H^2(M)$. The tangent map is

$$Z^1(M) \oplus C^0 \ni (x, c) \mapsto e^c * x = x - dc \in Z^1(M) \oplus d(C^0) = Z^1(M) \oplus D^1 = Z^1(L)$$

and it is an isomorphism. The inclusion $M \hookrightarrow L$ is a quasiisomorphism, therefore the obstruction to lifting x in M is equal to the obstruction to lifting $x = e^0 * x$ in L. We conclude the proof by observing that, according to [4, Prop. 7.5], [19, Lemma 2.21], the obstruction maps of Maurer-Cartan functor are invariant under the gauge action. □

Corollary A.1. *Let M be a differential graded Lie algebra, $L = M[t, dt]$ and $C \subseteq M[t]$ the subspace consisting of polynomials $g(t)$ with $g(0) = 0$. Then for every $A \in$ **Art** the map $(x, g[t]) \mapsto e^{g(t)} * x$ induces an isomorphism*

$$\mathrm{MC}_M(A) \times (C^0 \otimes \mathfrak{m}_A) \simeq \mathrm{MC}_L(A).$$

Proof. The data M, C and $D = d(C)$ satisfy the condition of Proposition A.1. □

Corollary A.2. *Let M be a differential graded Lie algebra. Two elements $x_0, x_1 \in \mathrm{MC}_M(A)$ are gauge equivalent if and only if they are homotopy equivalent.*

Proof. If x_0 and x_1 are gauge equivalent, then there exists $g \in M^0 \otimes \mathfrak{m}_A$ such that $e^g * x_0 = x_1$. Then, by Corollary A.1. $x(t) = e^{t\,g} * x_0$ is an element of $\mathrm{MC}_{M[t,dt]}(A)$ with $x(0) = x_0$ and $x(1) = x_1$, i.e., x_0 and x_1 are homotopy equivalent.
Vice versa, if x_0 and x_1 are homotopy equivalent, there exists $x(t) \in \mathrm{MC}_{M[t,dt]}(A)$ such that $x(0) = x_0$ and $x(1) = x_1$. By Corollary A.1., there exists $g(t) \in M^0[t] \otimes \mathfrak{m}_A$ with $g(0) = 0$ such that $x(t) = e^{g(t)} * x_0$. Then $x_1 = e^{g(1)} * x_0$, i.e., x_0 and x_1 are gauge equivalent. □

Part 2. dg-Grassmann functors

Let W be vector space over $\mathbb{K}$. The total Grassmannian of W is

$$\mathrm{Grass}(W) = \{\text{linear subspaces of } W\}.$$

The group $\mathrm{Aut}(W)$ of linear automorphisms of W acts on $\mathrm{Grass}(W)$. Denoting by $\mathrm{Grass}(V, W) \subseteq \mathrm{Grass}(W)$ the orbit of a subspace $V \subseteq W$ we have

$$\mathrm{Grass}(V, W) = \frac{\mathrm{Aut}(W)}{\mathrm{Aut}(V, W)}, \quad \text{where} \quad \mathrm{Aut}(V, W) = \{g \in \mathrm{Aut}(W) \mid g(V) = V\}.$$

The infinitesimal neighborhood of V in $\operatorname{Grass}(W)$ is the formal moduli space for the functor $\operatorname{Grass}_{V,W}\colon \mathbf{Art}\to\mathbf{Set}$:

$$\begin{aligned}\operatorname{Grass}_{V,W}(A) =&\{\phi\colon \operatorname{Spec}(A)\to\operatorname{Grass}(W)\mid \phi(\operatorname{Spec}(\mathbb{K}))=V\}\\ =&\{\text{free } A \text{ submodules } V_A\subseteq W\otimes A\mid V_A\otimes_A\mathbb{K}=V\}\\ =&\{f(V\otimes A)\subseteq W\otimes A\mid f\in\operatorname{Aut}(W\otimes A),\ f_{|W}=\operatorname{Id}\}.\end{aligned}$$

Since $\{f\in\operatorname{Aut}(W\otimes A)\mid f_{|W}=\operatorname{Id}\}=\exp(\mathfrak{gl}(W)\otimes\mathfrak{m}_A)$ we can write

$$\operatorname{Grass}_{V,W}(A)=\frac{\exp(\mathfrak{gl}(W)\otimes\mathfrak{m}_A)}{\exp(L^0_{V,W}\otimes\mathfrak{m}_A)}$$

where

$$L^0_{V,W}=\{g\in\mathfrak{gl}(W)\mid g(V)\subseteq V\}$$

and the action is given by

$$\exp(L^0_{V,W}\otimes\mathfrak{m}_A)\times\exp(\mathfrak{gl}(W)\otimes\mathfrak{m}_A)\to\exp(\mathfrak{gl}(W)\otimes\mathfrak{m}_A),\qquad (e^a,e^m)\mapsto e^me^{-a}.$$

In conclusion, the functor $\operatorname{Grass}_{V,W}$ coincides with the deformation functor Def_χ where $\chi\colon L^0_{V,W}\to\mathfrak{gl}(W)$ is the inclusion. Indeed the differential graded Lie algebras $L^0_{V,W}$ and $\mathfrak{gl}(W)$ are concentrated in degree 0, $\operatorname{MC}_\chi(A)=\exp(\mathfrak{gl}(W)\otimes\mathfrak{m}_A)$ and the gauge action is given by

$$\exp(L^0_{V,W}\otimes\mathfrak{m}_A)\times\exp(\mathfrak{gl}(W)\otimes\mathfrak{m}_A)\to\exp(\mathfrak{gl}(W)\otimes\mathfrak{m}_A),\qquad (e^a,e^m)\mapsto e^me^{-\chi(a)}.$$

8. The coarse dg-Grassmannian

The considerations of the above section suggest the following generalization from vector spaces to differential complexes. Let (W,d) be a dg-vector space and denote by:

(1) $\operatorname{Aut}(W)$ the group of automorphisms of the graded vector space W;
(2) $\operatorname{Aut}(W,d)$ the group of automorphisms of the differential graded vector space (W,d), i.e. the subgroup of $\operatorname{Aut}(W,d)$ consisting of linear automorphisms which commute with the differential d;
(3) $\operatorname{Aut}^0(W,d)$ the subgroup of $\operatorname{Aut}(W,d)$ of automorphisms inducing the identity in cohomology.

Define the coarse Grassmannian $\operatorname{Grass}(W)$ as the quotient

$$\operatorname{Grass}(W)=\frac{G(W)}{\operatorname{Aut}^0(W,d)},\qquad\text{where}\qquad G(W)=\{\text{subcomplexes of }(W,d)\,\}.$$

Note that $\operatorname{Aut}(W,d)$ acts on $G(W)$ and then the quotient group $\operatorname{Aut}(W,d)/\operatorname{Aut}^0(W,d)$ acts on $\operatorname{Grass}(W)$.

If $d=0$ then $\operatorname{Aut}^0(W,d)=\{\operatorname{Id}\}$ and then $\operatorname{Grass}(W)$ is the standard Grassmannian. Notice that the cohomology functor gives a map

$$h\colon \operatorname{Grass}(W)\to\operatorname{Grass}(H^*(W)),\qquad h(V)=\operatorname{Im}(H^*(V)\to H^*(W)).$$

Denoting by $\operatorname{Grass}(W)^s\subseteq\operatorname{Grass}(W)$ the “open” subset consisting of subcomplexes $V\subseteq W$ such that $H^*(V)\to H^*(W)$ is injective, we shall prove later (Theorems 9.3 and 10.6) that, is some sense, the map $h\colon \operatorname{Grass}(W)^s\to\operatorname{Grass}(H^*(W))$ is a local isomorphism; this fact justifies the quotient of $G(W)$ by the action of $\operatorname{Aut}^0(W,d)$.

Given a subcomplex $V \subseteq W$ we denote $\operatorname{Grass}(V,W) = G(V,W)/\operatorname{Aut}^0(W)$, where $G(V,W)$ the set of subcomplexes of W that are isomorphic to V as graded vector spaces; equivalently $U \in G(V,W)$ if and only if $U \in G(W)$ and there exists $f \in \operatorname{Aut}(W)$ such that $f(V) = U$ and then we have a natural identification

$$\frac{\{f \in \operatorname{Aut}(W) \mid df(V) \subseteq f(V)\}}{\operatorname{Aut}(V,W)} \xrightarrow{\simeq} G(V,W), \qquad f \mapsto f(V),$$

where $\operatorname{Aut}(V,W)$ is the subgroup of $\operatorname{Aut}(W)$ consisting of the automorphisms g such that $g(V) = V$.

Remark 8.1. To put some structure on $\operatorname{Grass}(W)$ the natural choice is to take $\operatorname{Grass}(V,W)$ as its components; then try to define $\operatorname{Grass}(V,W)$ as categorical quotient (in a suitable category: varieties, schemes, stacks, ...) using the identification

$$\operatorname{Grass}(V,W) = \frac{\{f \in \operatorname{Aut}(W) \mid df(V) \subseteq f(V)\}}{\operatorname{Aut}^0(W,d) \times \operatorname{Aut}(V,W)}.$$

Note that $(\varphi,\psi) \in \operatorname{Aut}^0(W,d) \times \operatorname{Aut}(V,W)$ acts of $\{f \in \operatorname{Aut}(W) \mid df(V) \subseteq f(V)\}$ by the formula $(\varphi,\psi)\cdot f = \varphi f \psi^{-1}$.
Unfortunately we may not expect that $\operatorname{Grass}(V,W)$ is separated in general. Consider $W^0 = W^1 = \mathbb{K} \oplus \mathbb{K}$, $W^i = 0$ for $i \neq 0,1$, and $d\colon W^0 \to W^1$ of rank 1.
If $V^0 = V^1 = \mathbb{K}$, then V is a subcomplex of W if and only if $V^0 = \ker(d)$ or $V^1 = \operatorname{Im}(d)$. Therefore $G(V,W) \subseteq \mathbb{P}(W^0) \times \mathbb{P}(W^1) = \mathbb{P}^1 \times \mathbb{P}^1$ is the union of two intersecting lines. In particular $G(V,W)$ is singular at the point $V^0 = \ker(d)$, $V^1 = \operatorname{Im}(d)$.
The group $\operatorname{Aut}^0(W,d)$ acts transitively on $\{V \mid V_0 \neq \ker(d)\}$, $\{V \mid V_1 \neq \operatorname{Im}(d)\}$ and then $\operatorname{Grass}(V,W)$ contains three point and it is not separated (same type of $\{xy = 0\}/\mathbb{K}^*$).

9. Infinitesimal study

Let (W,d) be a complex of vector spaces and $V \subseteq W$ a subcomplex; denote by L_W and $L_{V,W}$ the differential graded Lie algebras

$$L_W = \operatorname{Hom}^*(W,W), \qquad L_{V,W} = \{g \in \operatorname{Hom}^*(W,W) \mid g(V) \subseteq V\}.$$

An Artinian algebra can be seen as a differential complex with trivial differential; for every $A \in \mathbf{Art}$ we still denote by d the differential on $W \otimes A$. Since the differential on A is trivial, we have

$$d\colon W^i \otimes A \to W^{i+1} \otimes A, \qquad d(v \otimes a) = d(v) \otimes a.$$

Let $V \subseteq W$ be a subcomplex and consider the functors

$$\operatorname{Aut}_W, \operatorname{Aut}_{W,d}, \operatorname{Aut}^0_{W,d}, \operatorname{Aut}_{V,W}\colon \mathbf{Art} \to \mathbf{Groups}$$

defined as

$$\begin{aligned}
\operatorname{Aut}_W(A) &= \{f \in \operatorname{Hom}^0_A(W \otimes A, W \otimes A) \mid f \equiv \operatorname{Id} \pmod{\mathfrak{m}_A}\}\\
\operatorname{Aut}_{W,d}(A) &= \{f \in \operatorname{Aut}_W(A) \mid fd = df\}\\
\operatorname{Aut}_{V,W}(A) &= \{f \in \operatorname{Aut}_W(A) \mid f(V \otimes A) = V \otimes A\}\\
\operatorname{Aut}^0_{W,d}(A) &= \{f \in \operatorname{Aut}_{W,d}(A) \mid H^*(f) \text{ is the identity on } H^*(W \otimes A, d)\}
\end{aligned}$$

The above functors are smooth and homogeneous [29, 19]. Their tangent spaces are

$$\begin{aligned} T^1\operatorname{Aut}_W &= \operatorname{Hom}^0(W,W)=L^0_W\\ T^1\operatorname{Aut}_{W,d} &= \{f\in\operatorname{Hom}^0(W,W)\mid fd=df\}=Z^0(\operatorname{Hom}^*(W,W))\\ T^1\operatorname{Aut}_{W,V} &= \{f\in\operatorname{Hom}^0(W,W)\mid f(V)\subseteq V\}=L^0_{V,W}\\ T^1\operatorname{Aut}^0_{W,d} &= \{f\in\operatorname{Hom}^0(W,W)\mid fd=df,\ f(\ker(d))\subseteq\operatorname{Im}(d)\}\end{aligned}$$

Lemma 9.1. *For every $A\in\mathbf{Art}$, the exponential map gives isomorphisms*

$$\begin{aligned} &\exp\colon \operatorname{Hom}^0(W,W)\otimes\mathfrak{m}_A\to\operatorname{Aut}_W(A)\\ &\exp\colon Z^0(\operatorname{Hom}^*(W,W))\otimes\mathfrak{m}_A\to\operatorname{Aut}_{W,d}(A)\\ &\exp\colon B^0(\operatorname{Hom}^*(W,W))\otimes\mathfrak{m}_A\to\operatorname{Aut}^0_{W,d}(A)\\ &\exp\colon L^0_{V,W}\otimes\mathfrak{m}_A\to\operatorname{Aut}_{V,W}(A)\end{aligned}$$

Proof. First we note that For every i there exists a natural exact sequence

$$0\to B^i(\operatorname{Hom}^*(W,W))\to Z^i(\operatorname{Hom}^*(W,W))\to\operatorname{Hom}^i(H^*(W),H^*(W))\to 0$$

and therefore

$$T^1\operatorname{Aut}^0_{W,d}=B^0(\operatorname{Hom}^*(W,W))=\{df+fd\mid f\in\operatorname{Hom}^{-1}(W,W)\}.$$

Since all the functors are smooth and homogeneous, it is sufficient to prove that the exponential induces isomorphisms on the tangent spaces. □

The considerations made in Section 8 lead us to define the functors

$$M_{V,W},\operatorname{Grass}_{V,W}\colon\mathbf{Art}\to\mathbf{Sets}$$

$$M_{V,W}(A)=\{f\in\operatorname{Aut}_W(A)\mid df(V\otimes A)\subseteq f(V\otimes A)\},$$

$$\operatorname{Grass}_{V,W}=\frac{M_{V,W}}{\operatorname{Aut}^0_{W,d}\times\operatorname{Aut}_{V,W}}.$$

Proposition 9.2. *In the notation above, let $\chi\colon L_{V,W}\to L_W$ the inclusion. Then there exist natural isomorphisms of functors*

$$\operatorname{MC}_\chi\xrightarrow{\sim}M_{V,W},\qquad \operatorname{Def}_\chi\xrightarrow{\sim}\operatorname{Grass}_{V,W}.$$

Proof. Note that, since V is a subcomplex of (W,d), we have $d(V\otimes A)\subseteq V\otimes A$, and so

$$\begin{aligned} M_{V,W}(A)&=\{f\in\operatorname{Aut}_W(A)\mid (f^{-1}df-d)(V\otimes A)\subseteq V\otimes A\}=\\ &=\{f\in\operatorname{Aut}_W(A)\mid f^{-1}df-d\in L^1_{V,W}\otimes\mathfrak{m}_A\},\end{aligned}$$

and then, by Lemma 9.1 and by the identity $e^{-a}*0=e^{-a}de^a-d$,

$$M_{V,W}(A)=\{e^a\in\exp(L^0_W\otimes\mathfrak{m}_A)\mid e^{-a}*0\in L^1_{V,W}\otimes\mathfrak{m}_A\}.$$

Recall that

$$\operatorname{MC}_\chi(A)=\left\{(x,e^a)\in(L^1_{V,W}\otimes\mathfrak{m}_A)\times\exp(L^0_W\otimes\mathfrak{m}_A)\mid dx+\frac{1}{2}[x,x]=0,\ e^{-a}*0=\chi(x)\right\}.$$

Since χ is injective and the set of solutions of the Maurer-Cartan equation in $L^1_W\otimes\mathfrak{m}_A$ is preserved by the gauge action $l\mapsto e^{-a}*l$, we have

$$\operatorname{MC}_\chi(A)=\left\{e^a\in\exp(L^0_W\otimes\mathfrak{m}_A)\mid e^{-a}*0\in L^1_{V,W}\otimes\mathfrak{m}_A\right\}$$

and then the isomorphism $\mathrm{MC}_\chi \simeq M_{V,W}$.
The gauge action

$$(\exp(L^0_{V,W} \otimes \mathfrak{m}_A) \times \exp(B^0(L_W \otimes \mathfrak{m}_A))) \times \mathrm{MC}_\chi(A) \xrightarrow{*} \mathrm{MC}_\chi(A)$$

becomes

$$(e^l, e^{dm}) * e^a = e^{dm} e^a e^{-\chi(l)} = e^{dm} e^a e^{-l}.$$

According to Lemma 9.1 we have

$$\exp(L^0_{V,W} \otimes \mathfrak{m}_A) \times \exp(B^0(L_W \otimes \mathfrak{m}_A)) = \mathrm{Aut}_{V,W}(A) \times \mathrm{Aut}^0_{W,d}(A)$$

and it is immediate to observe that the gauge action is identified with the natural group action

$$\mathrm{Aut}_{V,W}(A) \times \mathrm{Aut}^0_{W,d}(A) \times M_{V,W}(A) \to M_{V,W}(A)$$

and then we have the isomorphism of quotient functors

$$\mathrm{Grass}_{V,W} = \mathrm{Def}_\chi\,.$$

□

Note that the isomorphism $\mathrm{Def}_\chi \to \mathrm{Grass}_{V,W}$ has a natural explicit description: it is the map

$$e^a \mapsto (e^a(V \otimes A), d) \subseteq (W \otimes A, d).$$

Theorem 9.3. *Let V be a subcomplex of (W, d) such that the inclusion $V \hookrightarrow W$ induces an injective morphism in cohomology $H^*(V) \hookrightarrow H^*(W)$. Identifying $H^*(V)$ with its image in $H^*(W)$, then the cohomology functor H^* gives a natural transformation of functors*

$$H^*\colon\ \mathrm{Grass}_{V,W} \to \mathrm{Grass}_{H^*(V),H^*(W)} \simeq \prod_i \mathrm{Grass}_{H^i(V),H^i(W)}\,.$$

Proof. Recall that

$$\mathrm{Grass}_{H^*(V),H^*(W)}(A) = \{\text{free } A \text{ submodules } F_A \subseteq H^*(W) \otimes A \mid F_A \otimes_A \mathbb{K} = H^*(V)\}.$$

Given $e^a \in \mathrm{MC}_\chi(A)$ denote by $d_a = e^{-a} d e^a \colon W \otimes A \to W \otimes A$; then $(V \otimes A, d_a)$ is a subcomplex of $(W \otimes A, d_a)$ and e^a is a morphism of complexes $(W \otimes A, d_a) \to (W \otimes A, d)$. By local flatness criteria, the cohomology of $(V \otimes A, d_a)$ is a free A-module and the map

$$H^*(e^a(V \otimes A), d) \to H^*(W \otimes A, d) \simeq H^*(W) \otimes A$$

is injective since it factors as

$$H^*(e^a(V \otimes A), d) \simeq H^*(V \otimes A, d_a) \xrightarrow{H^*(e^a)} H^*(W \otimes A, d)$$

where the isomorphism on the left is $H^*(e^{-a})$. Therefore the map

$$h\colon\ \mathrm{MC}_\chi(A) \to \mathrm{Grass}_{H^*(V),H^*(W)}(A),$$

$$h(e^a) = \text{image of the natural map } H^*(e^a(V \otimes A)) \to H^*(W) \otimes A$$

is well defined and factors to a natural transformation of functors

$$H^*\colon\ \mathrm{Grass}_{V,W} \longrightarrow \mathrm{Grass}_{H^*(V),H^*(W)}\,.$$

□

10. Homotopy invariance

Lemma 10.1. *If $V \hookrightarrow W$ is a quasiisomorphism, then the inclusion*

$$L_{V,W} \hookrightarrow L_W.$$

and the projection

$$L_{V,W} \to L_V, \qquad f \mapsto f|_V$$

are quasiisomorphisms of DGLA.

Proof. We have two short exact sequences

$$0 \to L_{V,W} \to L_W \to \operatorname{Hom}^*(V, W/V) \to 0$$

$$0 \to \operatorname{Hom}^*(W/V, W) \to L_{V,W} \to L_V \to 0$$

Since the complex W/V is acyclic and the bifunctor Hom^* commutes with cohomology, the complexes $\operatorname{Hom}^*(V, W/V)$ and $\operatorname{Hom}^*(V, W/V)$ are acyclic. □

Lemma 10.2. *Assume we have an exact diagram of differential graded vector spaces*

$$\begin{tikzcd}
 & & 0 \arrow[d] & & \\
 & & A \arrow[d, "\alpha"] & & \\
0 \arrow[r] & B \arrow[r, "\varphi"] & C \arrow[r, "\psi"] \arrow[d, "\beta"] & D \arrow[r] & 0 \\
 & & E \arrow[d] & & \\
 & & 0 & &
\end{tikzcd}$$

If $\psi\alpha$ is a quasiisomorphism, then also $\beta\varphi$ is a quasiisomorphism.

Proof. For every i in $\mathbb{Z}$ the map $H^i(\psi\alpha)\colon H^i(A) \to H^i(D)$ is an isomorphism. Since $H^i(\psi\alpha) = H^i(\psi) \circ H^i(\alpha)$, the map $H^i(\alpha)\colon H^i(A) \to H^i(C)$ is injective and $H^i(\psi)\colon H^i(C) \to H^i(D)$ is surjective. Therefore, the long exact sequences

$$\cdots \to H^{i-1}(E) \xrightarrow{\delta^{i-1}} H^i(A) \xrightarrow{H^i(\alpha)} H^i(C) \xrightarrow{H^i(\beta)} H^i(E) \xrightarrow{\delta^i} H^{i+1}(A) \to \cdots$$

and

$$\cdots \to H^{i-1}(D) \xrightarrow{\delta^{i-1}} H^i(B) \xrightarrow{H^i(\varphi)} H^i(C) \xrightarrow{H^i(\psi)} H^i(D) \xrightarrow{\delta^i} H^{i+1}(B) \to \cdots$$

can be refined as

$$H^{i-1}(E) \xrightarrow{\delta^{i-1}} 0 \to H^i(A) \xrightarrow{H^i(\alpha)} H^i(C) \xrightarrow{H^i(\beta)} H^i(E) \xrightarrow{\delta^i} 0$$

and

$$H^{i-1}(D) \xrightarrow{\delta^{i-1}} 0 \to H^i(B) \xrightarrow{H^i(\varphi)} H^i(C) \xrightarrow{H^i(\psi)} H^i(D) \xrightarrow{\delta^i} 0$$

and we have an exact diagram

$$\begin{tikzcd}
 & & 0 \arrow[d] & & \\
 & & H^i(A) \arrow[d, "H^i(\alpha)"] & & \\
0 \arrow[r] & H^i(B) \arrow[r, "H^i(\varphi)"] & H^i(C) \arrow[r, "H^i(\psi)"] \arrow[d, "H^i(\beta)"] & H^i(D) \arrow[r] & 0 \\
 & & H^i(E) \arrow[d] & & \\
 & & 0 & &
\end{tikzcd}$$

Now it is an easy exercise on linear algebra to show that $H^i(\beta\varphi) = H^i(\beta) \circ H^i(\varphi)$ is an isomorphism, i.e., $\beta\varphi$ is a quasiisomorphism. Note that, by symmetry, if $\beta\varphi$ is a quasiisomorphism, then also $\psi\alpha$ is a quasiisomorphism. $\square$

Lemma 10.3. *Consider a subcomplex U of $V \subseteq W$ and denote*

$$L_{U,V,W} = L_{U,W} \cap L_{V,W} = \{f \in \mathrm{Hom}^*(W,W) \mid f(U) \subseteq U,\ f(V) \subseteq V\}.$$

1. *If $U \hookrightarrow V$ is a quasiisomorphism, then the two inclusions*
$$L_{U,V,W} \hookrightarrow L_{V,W}, \qquad L_{U,V,W} \hookrightarrow L_{U,W}$$
are quasiisomorphisms of DGLA.
2. *If $V \hookrightarrow W$ is a quasiisomorphism, then the inclusions $L_{U,V,W} \hookrightarrow L_{U,W}$ and the projection*
$$L_{U,V,W} \to L_{U,V}, \qquad f \mapsto f_{|V}$$
are quasiisomorphisms of DGLA.

Proof. Assume that $U \hookrightarrow V$ is a quasiisomorphism; consider the exact sequences

$$0 \to L_{U,V,W} \xrightarrow{(\mathrm{id}|_V,\mathrm{id})} L_{U,V} \oplus L_{V,W} \xrightarrow{\tau} L_V \to 0, \qquad \tau(f,g) = f - g_{|V}$$

and

$$0 \to L_{U,V} \xrightarrow{(\mathrm{id},0)} L_{U,V} \oplus L_{V,W} \xrightarrow{\pi_2} L_{V,W} \to 0.$$

The composition $\tau \circ (\mathrm{id}, 0)\colon L_{U,V} \to L_V$ coincides with the inclusion $L_{U,V} \hookrightarrow L_V$. By Lemma 10.1, the inclusion $L_{U,V} \hookrightarrow L_V$ is a quasiisomorphism. Therefore, by Lemma 10.2 the map $\pi_2 \circ (\mathrm{id}|_V, \mathrm{id})\colon L_{U,V,W} \to L_{V,W}$ is a quasiisomorphism, i.e., $L_{U,V,W} \hookrightarrow L_{V,W}$ is a quasiisomorphism.

The acyclicity of V/U, together with the exact sequence

$$0 \to L_{U,V,W} \to L_{U,W} \xrightarrow{\sigma} \mathrm{Hom}^*\left(\frac{V}{U}, \frac{W}{V}\right) \to 0, \qquad \sigma(g) = g_{|V},$$

imply that $L_{U,V,W} \hookrightarrow L_{U,W}$ is a quasiisomorphism.
The proof of the second statement is very similar. Assume that $V \hookrightarrow W$ is a quasiisomorphism and consider the exact sequences

$$0 \to L_{U,V,W} \xrightarrow{(\mathrm{id}|_V,\mathrm{id})} L_{U,V} \oplus L_{V,W} \xrightarrow{\tau} L_V \to 0, \qquad \tau(f,g) = g_{|V} - f$$

and

$$0 \to L_{V,W} \xrightarrow{(0,\mathrm{id})} L_{U,V} \oplus L_{V,W} \xrightarrow{\pi_1} L_{U,V} \to 0.$$

The projection $L_{V,W} \to L_V$ is a quasiisomorphism by Lemma 10.1. This projection can be written as the composition $\tau \circ (0, \mathrm{id})\colon L_{V,W} \to L_V$, hence by Lemma 10.2 the map $\pi_1 \circ (\mathrm{id}|_V, \mathrm{id})\colon L_{U,V,W} \to L_{U,V}$ is a quasiisomorphism, i.e., the projection $L_{U,V,W} \to L_{U,V}$ is a quasiisomorphism. Moreover W/V is acyclic and then the inclusion $L_{U,V,W} \hookrightarrow L_{U,W}$ is a quasiisomorphism. $\square$

Lemma 10.4. *Assume we have a commutative diagram of* ***inclusions*** *of differential graded vector spaces*

$$\begin{tikzcd} U \arrow[r] \arrow[d] & V \arrow[d] \\ Z \arrow[r] & W \end{tikzcd}$$

If the horizontal arrows are quasiisomorphisms then $\mathrm{Grass}_{U,Z} \simeq \mathrm{Grass}_{U,W} \simeq \mathrm{Grass}_{V,W}$.

Proof. In the notation of Lemma 10.3, there exists a commutative diagram of DGLA, where every arrow is either the natural inclusion or the natural projection:

$$\begin{tikzcd} L_{U,Z} \arrow[d, "\eta"] & L_{U,Z,W} \arrow[l] \arrow[r] \arrow[d] & L_{U,W} \arrow[d, "\rho"] & L_{U,V,W} \arrow[l] \arrow[r] \arrow[d] & L_{V,W} \arrow[d, "\chi"] \\ L_Z & L_{Z,W} \arrow[l] \arrow[r] & L_W & L_W \arrow[l] \arrow[r] & L_W \end{tikzcd}$$

By Lemma 10.3 the horizontal arrows are quasiisomorphism and then, Theorem 6.1 gives an isomorphism of deformation functors

$$\mathrm{Grass}_{U,Z} \simeq \mathrm{Def}_\eta \simeq \mathrm{Def}_\rho \simeq \mathrm{Grass}_{U,W} \simeq \mathrm{Def}_\chi \simeq \mathrm{Grass}_{V,W}\,.$$

$\square$

Remark 10.5. In the same hypothesis of Lemma 10.4, if in addition $U = V \cap Z$, then the isomorphism $\mathrm{Grass}_{U,Z} \simeq \mathrm{Grass}_{V,W}$ can be done in a more explicit and easy way. In fact the two inclusion

$$\alpha\colon L_{Z,W} \cap L_{V,W} \hookrightarrow L_{U,V,W}, \qquad \beta\colon L_{Z,W} \cap L_{V,W} \hookrightarrow L_{U,Z,W}$$

are quasiisomorphisms since their cokernels are the acyclic complexes

$$\mathrm{Coker}(\alpha) = \mathrm{Hom}^*\left(\frac{Z}{U}, \frac{W}{Z}\right), \qquad \mathrm{Coker}(\beta) = \mathrm{Hom}^*\left(\frac{V}{U}, \frac{W}{V}\right).$$

Then, according to Lemma 10.4 every horizontal arrow of

$$\begin{tikzcd} L_{U,Z} \arrow[d, "\eta"] & L_{Z,W} \cap L_{V,W} \arrow[l] \arrow[r, hook] \arrow[d] & L_{V,W} \arrow[d, "\chi"] \\ L_Z & L_{Z,W} \arrow[l] \arrow[r, hook] & L_W \end{tikzcd}$$

is a quasiisomorphism. The isomorphism $\mathrm{Grass}_{V,W} \simeq \mathrm{Grass}_{U,Z}$ can then be explicitly described as follows: given $A \in \mathbf{Art}$ and an element $e^a(V \otimes A)$ in $\mathrm{Grass}_{V,W}(A)$, there exists (and it is unique up to gauge) an element $e^\alpha \in \exp(\mathrm{Hom}^0(W, W) \otimes \mathfrak{m}_A)$ such that $e^\alpha(V \otimes A) = e^a(V \otimes A)$ and $e^\alpha(Z \otimes A) \subseteq Z \otimes A$. Then the isomorphism $\mathrm{Grass}_{V,W} \simeq \mathrm{Grass}_{U,Z}$ is given by the map

$$e^a(V \otimes A) \mapsto e^\alpha(U \otimes A) \subseteq Z \otimes A.$$

Theorem 10.6. *Let V be a subcomplex of (W, d) such that the inclusion $V \hookrightarrow W$ induces an injective morphism in cohomology $H^*(V) \hookrightarrow H^*(W)$. Identifying $H^*(V)$*

with its image in $H^*(W)$*, then the cohomology functor* H^* *gives an isomorphism of functors*

$$H^*\colon \operatorname{Grass}_{V,W} \xrightarrow{\sim} \operatorname{Grass}_{H^*(V),H^*(W)} \cong \prod_i \operatorname{Grass}_{H^i(V),H^i(W)}.$$

Proof. It is possible to find "harmonic representatives" $\mathcal{H}_V \subseteq V$ and $\mathcal{H}_W \subseteq W$ with $\mathcal{H}_V = \mathcal{H}_W \cap V$. Indeed, the injectivity of $H^*(V) \to H^*(W)$ implies the two equalities

$$Z^*(V) = V \cap Z^*(W), \qquad B^*(V) = V \cap B^*(W).$$

Therefore it is possible to find a subspace $\mathcal{H}_W \subseteq Z^*(W)$ such that $\mathcal{H}_W \oplus B^*(W) = Z^*(W)$ and $(\mathcal{H}_W \cap V) \oplus B^*(V) = Z^*(V)$. Set $\mathcal{H}_V = \mathcal{H}_W \cap V$. Then the diagram

$$\begin{tikzcd}
\mathcal{H}_V \arrow[r] \arrow[d] & V \arrow[d] \\
\mathcal{H}_W \arrow[r] & W
\end{tikzcd}$$

satisfies the hypothesis of Lemma 10.4 and Remark 10.5 and we have an isomorphism $\operatorname{Grass}_{V,W} \xrightarrow{\sim} \operatorname{Grass}_{\mathcal{H}_V,\mathcal{H}_W}$ given by

$$e^a(V \otimes A) \mapsto e^\alpha(\mathcal{H}_V \otimes A),$$

where $e^\alpha \in \exp(\operatorname{Hom}^0(W,W) \otimes \mathfrak{m}_A)$ is such that $e^\alpha(V \otimes A) = e^a(V \otimes A)$ and $e^\alpha(\mathcal{H}_W \otimes A) \subseteq \mathcal{H}_W \otimes A$.

Now, according to Theorem 9.3 it is sufficient to note that the natural map

$$e^\alpha(\mathcal{H}_V \otimes A) \to H^*(e^\alpha(V \otimes A), d) = H^*(e^a(V \otimes A), d)$$

is an isomorphism. □

Part 3. The universal period map

In this part we work over the field $\mathbb{K} = \mathbb{C}$ of complex numbers. Unless otherwise specified, the symbol $\otimes$ denotes tensor product over $\mathbb{C}$.

11. The Kodaira-Spencer DGLA

We will follows the same general notation of [33]; in particular, for a differentiable manifold M we denote by $\mathcal{A}^p_M$ the sheaf of differentiable p-forms with complex coefficients and by $d\colon \mathcal{A}^p_M \to \mathcal{A}^{p+1}_M$ the De Rham differential.

We think an almost complex structure on M as a subsheaf $\mathcal{V} \subseteq \mathcal{A}^1_M$ of locally free $\mathcal{A}^0_M$-modules such that $\mathcal{V} \oplus \overline{\mathcal{V}} = \mathcal{A}^1_M$.

For a complex manifold X we denote by:

- $T_{X,\mathbb{C}} = T^{1,0}_X \oplus T^{0,1}_X$ the complexified differential tangent bundle.
- $T_X \simeq T^{1,0}_X$ the holomorphic tangent bundle.
- $\mathcal{A}^{p,q}_X$ the sheaf of differentiable (p,q)-forms and by $\mathcal{A}^{p,q}_X(T_X)$ the sheaf of (p,q)-forms with values in T_X.
- $A^{p,q}_X$ and $A^{p,q}_X(T_X)$ the vector spaces of global sections of $\mathcal{A}^{p,q}_X$ and $\mathcal{A}^{p,q}_X(T_X)$ respectively.

Recall that an almost complex structure $\mathcal{V}\subseteq\mathcal{A}^1_X$ is called integrable if there exists a structure of complex manifold on X such that $\mathcal{V}=\mathcal{A}^{1,0}_X$.

The direct sum

$$\mathcal{A}_X=\mathop{\oplus}_{i}\mathcal{A}^i_X,\qquad\text{where}\qquad \mathcal{A}^i_X=\mathop{\oplus}_{p+q=i}\mathcal{A}^{p,q}_X,$$

endowed with the wedge product $\wedge$, is a sheaf of graded algebras; we denote by $\mathcal{D}er^{a,b}(\mathcal{A}_X)$ the sheaf of its $\mathbb{C}$-linear derivations of bidegree (a,b). Notice that ∂ and $\overline{\partial}$ are global sections of $\mathcal{D}er^{1,0}(\mathcal{A}_X)$ and $\mathcal{D}er^{0,1}(\mathcal{A}_X)$ respectively.
The direct sum $\mathcal{D}er^*(\mathcal{A}_X)=\oplus_k\oplus_{a+b=k}\mathcal{D}er^{a,b}(\mathcal{A}_X)$ is a sheaf of differential graded Lie algebras, with its natural bracket

$$[f,g]=fg-(-1)^{\deg(f)\deg(g)}gf$$

and differential $[d,-]=[\partial+\overline{\partial},-]$.
Similarly $\mathcal{A}^{0,*}_X(T_X)$ is a sheaf of DGLA, where the bracket $[\ ,\ ]$ and the differential D are defined in local coordinates by the formulas:

$$D\left(\phi\frac{\partial}{\partial z_i}\right)=-\overline{\partial}(\phi)\frac{\partial}{\partial z_i},\qquad \phi\in\mathcal{A}^{0,*}_X,$$

$$\left[fd\overline{z}_I\frac{\partial}{\partial z_i},gd\overline{z}_J\frac{\partial}{\partial z_j}\right]=d\overline{z}_I\wedge d\overline{z}_J\left(f\frac{\partial g}{\partial z_i}\frac{\partial}{\partial z_j}-g\frac{\partial f}{\partial z_j}\frac{\partial}{\partial z_i}\right).$$

The contraction of differential forms with vector fields is used to define two injective morphisms of sheaves: the *contraction map*

$$\boldsymbol{i}\colon\mathcal{A}^{0,*}_X(T_X)\to\mathcal{D}er^*(\mathcal{A}_X)[-1],\qquad a\mapsto\boldsymbol{i}_a,\quad \boldsymbol{i}_a(\omega)=a\lrcorner\omega,$$

and the *holomorphic Lie derivative*

$$\boldsymbol{l}\colon\mathcal{A}^{0,*}_X(T_X)\to\mathcal{D}er^*(\mathcal{A}_X),\qquad a\mapsto\boldsymbol{l}_a=[\partial,\boldsymbol{i}_a],\quad \boldsymbol{l}_a(\omega)=\partial(a\lrcorner\omega)+(-1)^{\deg(a)}a\lrcorner\partial\omega.$$

Lemma 11.1. *In the notation above, for every $a,b\in\mathcal{A}^{0,*}_X(T_X)$ we have*

$$\boldsymbol{i}_{Da}=-[\overline{\partial},\boldsymbol{i}_a],\qquad \boldsymbol{i}_{[a,b]}=[\boldsymbol{i}_a,[\partial,\boldsymbol{i}_b]],\qquad [\boldsymbol{i}_a,\boldsymbol{i}_b]=0.$$

In particular, since $[\partial,\boldsymbol{i}_b]=[d,\boldsymbol{i}_b]+\boldsymbol{i}_{Db}$, the contraction map $\boldsymbol{i}$ is a Cartan homotopy and the holomorphic Lie derivative is a morphism of sheaves of differential graded Lie algebras.

Proof. Since locally $\mathcal{A}_X$ is generated as $\mathbb{C}$-algebra by $\mathcal{A}^{0,0}_X\oplus\mathcal{A}^{1,0}_X\oplus\mathcal{A}^{0,1}_X$, if $\min(a,b)\leq-2$ then $\mathcal{D}er^{a,b}(\mathcal{A}_X)=0$, while if $\min(a,b)\leq-1$ then every $h\in\mathcal{D}er^{a,b}(\mathcal{A}_X)$ is C^∞-linear. The elements of the third formula belong to $\mathcal{D}er^{-2,*}(\mathcal{A}_X)$ and therefore vanish. Every term of first two formulas belongs to $\mathcal{D}er^{-1,*}(\mathcal{A}_X)$ and then it is sufficient to check equalities of such derivations when applied to dz_i, where $z_1,\dots,z_n$ are local holomorphic coordinates. This is straightforward and it is left to the reader: see also Lemma 7 of [21]. □

The contraction map gives a natural isomorphism of vector spaces

$$\boldsymbol{i}\colon A^{0,1}_X(T_X)\to\operatorname{Hom}_{\mathcal{A}^0_X}(\mathcal{A}^{1,0}_X,\mathcal{A}^{0,1}_X).$$

For every sufficiently small $\xi\in A^{0,1}_X(T_X)$, the graph of $\boldsymbol{i}_\xi\in\operatorname{Hom}_{\mathcal{A}^0_X}(\mathcal{A}^{1,0}_X,\mathcal{A}^{0,1}_X)$ determines a variation of the almost complex structure of X given by

$$\mathcal{A}^{1,0}_\xi=\{\omega\in\mathcal{A}^1_X\mid\overline{\pi}(\omega)=\boldsymbol{i}_\xi\pi(\omega)\}=\{\omega\in\mathcal{A}^1_X\mid\overline{\pi}(\omega)=\xi\lrcorner\pi(\omega)\},$$

where $\pi\colon \mathcal{A}^1_X \to \mathcal{A}^{1,0}_X$ and $\overline{\pi}\colon \mathcal{A}^1_X \to \mathcal{A}^{0,1}_X$ are the projections. Then denote

$$\mathcal{A}^{0,1}_\xi = \overline{\mathcal{A}^{1,0}_\xi}, \qquad \mathcal{A}^{p,q}_\xi = \bigwedge^p \mathcal{A}^{1,0}_\xi \otimes \bigwedge^q \mathcal{A}^{0,1}_\xi.$$

The sheaf of ξ-holomorphic functions is by definition

$$\mathcal{O}_\xi = \{f \in \mathcal{A}^0_X \mid df \in \mathcal{A}^{1,0}_\xi\}$$

and then, according to the definition of $\mathcal{A}^{1,0}_\xi$, we have

$$\mathcal{O}_\xi = \{f \in \mathcal{A}^0_X \mid \overline{\partial} f = \xi \lrcorner \partial f\} = \{f \in \mathcal{A}^0_X \mid (\overline{\partial} + \boldsymbol{l}_\xi) f = 0\}.$$

Since $\boldsymbol{i}_\xi$ is a nilpotent derivation of degree 0 of the sheaf of graded algebras $\mathcal{A}_X$, its exponential

$$e^{\boldsymbol{i}_\xi} = \exp(\xi \lrcorner)\colon \mathcal{A}_X \to \mathcal{A}_X$$

is an isomorphism of graded algebras.

Lemma 11.2. *In the above notation* $e^{\boldsymbol{i}_\xi}(\mathcal{A}^{1,0}_X) = \mathcal{A}^{1,0}_\xi$.

Proof. For every $\omega \in \mathcal{A}^1_X$ we have $e^{\boldsymbol{i}_\xi}(\omega) = \omega + \xi \lrcorner \pi(\omega)$, therefore

$$\pi e^{\boldsymbol{i}_\xi}(\omega) = \pi(\omega), \qquad \overline{\pi} e^{\boldsymbol{i}_\xi}(\omega) = \overline{\pi}(\omega) + \xi \lrcorner \pi(\omega)$$

and then $e^{\boldsymbol{i}_\xi}(\omega) \in \mathcal{A}^{1,0}_\xi$ if and only if $\overline{\pi}(\omega) = 0$. □

The Newlander-Nirenberg theorem [24], [13, Thm. 5.5] implies (see e.g. [2, Lecture 1]) that the following four conditions are equivalent:

(1) The almost complex structure $\mathcal{A}^{1,0}_\xi$ is integrable.
(2) ξ is a solution of the Maurer-Cartan equation [13, Equation 5.86], [2, Equation 2.5]
$$D\xi + \frac{1}{2}[\xi, \xi] = 0.$$
(3) For every $x \in X$ there exist $f_1, \ldots, f_n \in \mathcal{O}_{\xi,x}$, $n = \dim X$, such that $df_1, \ldots, df_n$ are a basis of the $\mathcal{A}^0_{X,x}$-module $\mathcal{A}^{1,0}_{\xi,x}$.
(4) $d\mathcal{F}^1_\xi \subseteq \mathcal{F}^1_\xi$, where $\mathcal{F}^1_\xi = \oplus_{p\geq 1} \mathcal{A}^{p,q}_\xi = (\mathcal{A}^{1,0}_\xi) \subseteq \mathcal{A}_X$ is the graded ideal sheaf generated by $\mathcal{A}^{1,0}_\xi$.

Definition 11.3. The Kodaira-Spencer algebra of a complex manifold X is the differential graded Lie algebra $K_X = \oplus_i A^{0,i}_X(T_X)$ of global sections of $\oplus_i \mathcal{A}^{0,i}_X(T_X)$.

Denoting by $\operatorname{Def}_X\colon \mathbf{Art} \to \mathbf{Set}$ the functor of infinitesimal deformations of X:

$$\operatorname{Def}_X(B) = \frac{\text{deformations of } X \text{ over } \operatorname{Spec}(B)}{\sim}.$$

A deformation of X over $\operatorname{Spec}(B)$ may be interpreted as a morphism $\mathcal{O}_B \to \mathcal{O}_X$ of sheaves of B-algebras such that $\mathcal{O}_B$ is flat over B and the induced map $\mathcal{O}_B \otimes_B \mathbb{C} \to \mathcal{O}_X$ is an isomorphism.

The following result is well known [3], [8], [14, Ex. 3.4.1]: a detailed proof will also appear in [12]

Theorem 11.4. *There exists an isomorphism of functors*

$$\mathcal{O}\colon \operatorname{Def}_{K_X} \to \operatorname{Def}_X$$

defined in the following way: given a local Artinian $\mathbb{C}$-algebra B and a solution of the Maurer-Cartan equation $\xi \in A^{0,1}_X(T_X)\otimes \mathfrak{m}_B$ we set

$$\mathcal{O}_\xi = \ker(\mathcal{A}^{0,0}_X \otimes B \xrightarrow{\overline{\partial}+\boldsymbol{l}_\xi} \mathcal{A}^{0,1}_X \otimes B) = \{f \in \mathcal{A}^{0,0}_X \otimes B \mid \overline{\partial} f = \xi \lrcorner \partial f\}$$

and the map $\mathcal{O}_\xi \to \mathcal{O}_X$ is induced by the projection $\mathcal{A}^{0,0}_X \otimes B \to \mathcal{A}^{0,0}_X \otimes \mathbb{C} = \mathcal{A}^{0,0}_X$.

12. The period map

Let X be a fixed complex manifold: we shall denote by $A_X = F^0 \supseteq F^1 \supseteq \cdots$ the Hodge filtration of differential forms on X, i.e. for every $p \geq 0$

$$F^p = \bigoplus_{i\geq p}\bigoplus_{j} A^{i,j}_X.$$

Theorem 12.1. *Let p be a fixed nonnegative integer and consider the inclusion of differential graded Lie algebras*

$$L_{F^p,A_X} = \{f \in \operatorname{Hom}^*(A_X,A_X) \mid f(F^p)\subseteq F^p\} \xrightarrow{\chi} \operatorname{Hom}^*(A_X,A_X) = L_{A_X}.$$

Then the linear map

$$\mathfrak{p}^p\colon K_X \to C_\chi, \qquad \mathfrak{p}^p(\xi) = (\boldsymbol{l}_\xi, \boldsymbol{i}_\xi) = ([\partial, \boldsymbol{i}_\xi], \boldsymbol{i}_\xi)$$

is a linear L_∞-morphism.
In particular $\mathfrak{p}^p$ induces a natural transformation of functors:

$$\mathfrak{p}^p\colon \operatorname{Def}_X \to \operatorname{Def}_\chi = \operatorname{Grass}_{F^p,A_X}.$$

Proof. According to Lemma 11.1 and Proposition 7.4, the Lie derivative

$$K_X = A^{0,*}_X(T_X) \xrightarrow{\boldsymbol{l}} \operatorname{Hom}^*(A_X,A_X)$$

is a morphism of differential graded Lie algebras and the map

$$\tilde{\boldsymbol{i}}\colon K_X \to C_{\boldsymbol{l}} = K_X \oplus \operatorname{Hom}^*(A_X,A_X)[-1], \qquad \tilde{\boldsymbol{i}}(\xi) = (\xi, \boldsymbol{i}_\xi)$$

is a linear L_∞-morphism.
The morphism $\mathfrak{p}^p$ is the composition of $\tilde{\boldsymbol{i}}$ and the linear L_∞-morphism $C_{\boldsymbol{l}} \to C_\chi$ induced by the horizontal arrows of the following commutative diagram of differential graded Lie algebras

$$\begin{tikzcd}
A^{0,*}_X(T_X) \arrow[r, "\boldsymbol{l}"] \arrow[d, "\boldsymbol{l}"] & L_{F^p,A_X} \arrow[d, "\chi"] \\
\operatorname{Hom}^*(A_X,A_X) \arrow[r, "="] & L_{A_X}
\end{tikzcd}$$

□

Corollary 12.2. *If $\xi \in A^{0,1}_X(T_X)$ is a solution of the Maurer-Cartan equation, then in the associative algebra $\operatorname{Hom}^*(A_X,A_X)$ we have the equality*

$$e^{-\boldsymbol{i}_\xi} d e^{\boldsymbol{i}_\xi} = d + e^{-\boldsymbol{i}_\xi} * 0 = d + \boldsymbol{l}_\xi = d + [\partial, \boldsymbol{i}_\xi].$$

Proof. The first equality follows from the explicit description of the gauge action in the DGLA $\operatorname{Hom}^*(A_X, A_X)$. Taking $p = 0$ in Theorem 12.1, the pair $\mathfrak{p}^0(\xi) = (\boldsymbol{l}_\xi, \boldsymbol{i}_\xi)$ satisfies the Maurer-Cartan equation in C_χ and then $e^{\boldsymbol{i}_\xi} * \boldsymbol{l}_\xi = 0$.
Obviously the above equality can be also proved directly as a consequence of Cartan homotopy formulas (Lemma 11.1). □

Theorem 12.3. *In the same notation of Theorem 12.1, assume that X is a compact Kähler manifold and denote by*

$$\mathcal{P}^p\colon \operatorname{Def}_X \to \operatorname{Grass}_{H^*(F^p), H^*(A_X)}$$

the composition of the natural transformation $\mathfrak{p}^p\colon \operatorname{Def}_X \to \operatorname{Grass}_{F^p, A_X}$ and the cohomology isomorphism $H^*\colon \operatorname{Grass}_{F^p, A_X} \to \operatorname{Grass}_{H^*(F^p), H^*(A_X)}$ (see Theorem 10.6). Then $\mathcal{P}^p$ is the universal period map.

Proof. Let $B \in \mathbf{Art}$ and $\xi \in \operatorname{MC}_{K_X}(B)$; by definition

$$\mathcal{P}^p(\xi) = H^*(e^{\boldsymbol{i}_\xi}(F^p \otimes B)) \subseteq H^*(A_X \otimes B) = H^*(A_X) \otimes B.$$

On the other hand, the period of the infinitesimal deformation $\mathcal{O}_\xi = \ker(\overline{\partial} + \boldsymbol{l}_\xi)$ is the B-submodule $H^*(F^p_\xi) \subseteq H^*(A_X \otimes B)$, where F^p_ξ is the complex of global sections of the differential ideal sheaf $\mathcal{F}^p_\xi \subseteq \mathcal{A}_X \otimes B$ generated by $(d\mathcal{O}_\xi)^p$.
It is sufficient to prove that $e^{\boldsymbol{i}_\xi}(F^p \otimes B) = F^p_\xi$; since $e^{\boldsymbol{i}_\xi}\colon \mathcal{A}_X \otimes B \to \mathcal{A}_X \otimes B$ is a morphism of sheaves of B-algebras, it is sufficient to prove that $e^{-\boldsymbol{i}_\xi}(d\mathcal{O}_\xi) \subseteq \mathcal{A}^{1,0}_X \otimes B$. This equality, together rank considerations, will imply that

$$e^{-\boldsymbol{i}_\xi}(\mathcal{F}^p_\xi) = \bigoplus_{i \geq p, j} \mathcal{A}^{i,j}_X \otimes B.$$

Since $e^{\boldsymbol{i}_\xi}$ is the identity on $\mathcal{A}^{0,0}_X \otimes B$, by Corollary 12.2 we can write

$$e^{-\boldsymbol{i}_\xi}(d\mathcal{O}_\xi) = e^{-\boldsymbol{i}_\xi} d e^{\boldsymbol{i}_\xi} \mathcal{O}_\xi = (\partial + \overline{\partial} + \boldsymbol{l}_\xi)\mathcal{O}_\xi = \partial\mathcal{O}_\xi \subseteq \partial\mathcal{A}^{0,0}_X \otimes B \subseteq \mathcal{A}^{1,0}_X \otimes B.$$

□

Remark 12.4. In the statement of Theorem 12.3 the Kähler assumption is used in cohomological sense; more precisely we only require that the cohomology of the complex F^p injects into the De Rham cohomology of X.

Corollary 12.5 (Griffiths)**.** *The differential of the universal period map is*

$$d\mathcal{P}^p = \boldsymbol{i}\colon H^1(X, T_X) \to \bigoplus_i \operatorname{Hom}\left(F^p H^i(X, \mathbb{C}), \frac{H^i(X, \mathbb{C})}{F^p H^i(X, \mathbb{C})}\right).$$

Proof. Recall from Remark 3.1 that the projection on the second factor induces an identification

$$H^j(C_\chi) \xrightarrow{\simeq} H^{j-1}(\operatorname{Hom}^*(F^p, A_X/F^p)) = \bigoplus_i \operatorname{Hom}(H^i(F^p), H^{i+j-1}(A_X/F^p)).$$

Via this identification, the L_∞-morphism $\mathfrak{p}^p\colon K_X \to C_\chi$, induces in cohomology the map

$$H^j(\mathfrak{p}^p) = \boldsymbol{i}\colon H^j(K_X) \to \bigoplus_i \operatorname{Hom}(H^i(F^p), H^{i+j-1}(A_X/F^p)).$$

The differential of $\mathcal{P}^p\colon \operatorname{Def}_X \to \operatorname{Grass}_{F^p, A_X} \simeq \operatorname{Grass}_{H^*(F^p), H^*(X;\mathbb{C})}$ is therefore

$$d\mathcal{P}^p = \boldsymbol{i}\colon H^1(K_X) \to \bigoplus_i \operatorname{Hom}(H^i(F^p), H^i(A_X/F^p)).$$

□

Corollary 12.6 (Kodaira's Principle [3, 21, 26])**.** *The obstructions to deformations of X are contained in the kernel of*

$$\boldsymbol{i}\colon H^2(X,T_X)\to\bigoplus_i \operatorname{Hom}\left(F^pH^i(X,\mathbb{C}),\frac{H^{i+1}(X,\mathbb{C})}{F^pH^{i+1}(X,\mathbb{C})}\right),$$

for every $p\geq 0$.

Proof. We use the same general argument of [21, Section 5]: since the period map $\mathcal{P}^p\colon \operatorname{Def}_X\to\operatorname{Grass}_{H^*(F^p),H^*(X,\mathbb{C})}$ is induced by the L_∞-morphism $\mathfrak{p}^p\colon K_X\to C_\chi$, the linear map

$$H^2(\mathcal{P}^p)\colon H^2(X,T_X)\to H^2(C_\chi)$$

is a morphism of obstruction theories, i.e., it commutes with the natural obstruction maps for Def_X and $\operatorname{Grass}_{H^*(F^p),H^*(X,\mathbb{C})}$ [23]. In particular, obstructions to deforming the complex structure of the Kähler manifold X are mapped to obstructions of the functor $\operatorname{Grass}_{H^*(F^p),H^*(X,\mathbb{C})}$. Since the latter is unobstructed, the obstructions to deforming X are annihilated by $H^2(\mathfrak{p}^p)$. By the proof of Corollary 12.5, $H^2(\mathfrak{p}^p)=\boldsymbol{i}$. □

13. Trasversality

Consider a fixed compact Kähler manifold X and a differential graded commutative $\mathbb{C}$-algebra (Ω,d_Ω). Let

$$\operatorname{Id}\otimes d_\Omega\colon A_X\otimes\Omega\to A_X\otimes\Omega,\qquad(\operatorname{Id}\otimes d_\Omega)(a\otimes\omega)=(-1)^{\deg(a)}a\otimes d_\Omega(\omega)$$

the trivial extension of d_Ω. Then $\operatorname{Id}\otimes d_\Omega$ is a differential of the graded algebra $A_X\otimes\Omega$ inducing a flat connection

$$\operatorname{Id}\otimes d_\Omega\colon H^*(A_X)\otimes\Omega^i\to H^*(A_X)\otimes\Omega^{i+1}.$$

Assume now that B is a commutative unital $\mathbb{C}$-algebra and let $\phi\colon B\to\Omega^0$ be a morphism of graded unital $\mathbb{C}$-algebras.
Via the natural isomorphisms

$$A_X\otimes\Omega=(A_X\otimes B)\otimes_B\Omega,\qquad H^*(A_X)\otimes\Omega=H^*(A_X\otimes B)\otimes_B\Omega$$

the operator $\operatorname{Id}\otimes d_\Omega$ induces the differential

$$\nabla\colon (A_X\otimes B)\otimes_B\Omega\to(A_X\otimes B)\otimes_B\Omega,$$
$$\nabla((a\otimes b)\otimes_B\omega)=(-1)^{\deg(a)}(a\otimes 1)\otimes_B d_\Omega(\phi(b)\omega)$$

and the flat connection

$$\nabla\colon H^*(A_X\otimes B)\otimes_B\Omega^i\to H^*(A_X\otimes B)\otimes_B\Omega^{i+1},$$

that, by analogy with the case of B a power series ring and $\Omega=\bigwedge_B^*\Omega_B^1$ we shall call *Gauss-Manin connection.*

Assume now that $B\in\mathbf{Art}$ and consider a deformation of X over $\operatorname{Spec}(B)$ determined by the Kuranishi data $\xi\in\operatorname{MC}_{K_X}(B)\subseteq A_X^{0,1}(T_X)\otimes\mathfrak{m}_B$.

The classical Griffiths' trasversality theorem [10], [33, Prop. 10.18] generalizes to the following result:

Proposition 13.1 (Trasversality)**.** *Let F_ξ^p be the complex of global sections of the differential ideal sheaf $\mathcal{F}_\xi^p\subseteq\mathcal{A}_X\otimes B$ generated by $(d\mathcal{O}_\xi)^p$. Then*

$$\nabla(H^*(F_\xi^p)\otimes_B\Omega^i)\subseteq H^*(F_\xi^{p-1})\otimes_B\Omega^{i+1}.$$

Proof. It is not restrictive to assume $B \subseteq \Omega^0 \subseteq \Omega$ and ϕ the inclusion. For notational simplicity denote by $\delta\colon A_X^{0,*}(T_X)\otimes\Omega \to A_X^{0,*}(T_X)\otimes\Omega$ the trivial extension of d_Ω, i.e. $\delta(\xi\otimes b) = (-1)^{\deg(\xi)}\xi\otimes d_\Omega(b)$.
It is sufficient to prove that $\nabla(F_\xi^p\otimes_B\Omega^i)\subseteq F_\xi^{p-1}\otimes_B\Omega^{i+1}$.

The contraction map $\boldsymbol{i}$ extends naturally to a Cartan homotopy (Example 7.3)

$$\boldsymbol{i}\colon A_X^{0,*}(T_X)\otimes\Omega\to \operatorname{Hom}^*(A_X,A_X)\otimes\Omega\subseteq \operatorname{Hom}^*((A_X\otimes B)\otimes_B\Omega,(A_X\otimes B)\otimes_B\Omega)$$

such that $\boldsymbol{i}_{\delta x} = -[\nabla,\boldsymbol{i}_x]$; in particular $[\boldsymbol{i}_x,[\boldsymbol{i}_x,\nabla]] = -[\boldsymbol{i}_x,\boldsymbol{i}_{\delta x}] = 0$ for every x. For every nilpotent $a\in \operatorname{Hom}^0((A_X\otimes B)\otimes_B\Omega,(A_X\otimes B)\otimes_B\Omega)$ we have the equality

$$\begin{aligned}[\nabla,e^a] &= \nabla e^a - e^a\nabla = e^a(e^{-a}\nabla e^a-\nabla)\\ &= e^a(e^{-\operatorname{ad}_a}(\nabla)-\nabla) = e^a[\nabla,a] - e^a\sum_{n\geq 2}\frac{(-\operatorname{ad}_a)^n}{n!}(\nabla).\end{aligned}$$

Since $\xi\in \mathrm{MC}_{K_X}(B)$, for $a\in A_X\otimes B$ and $\omega\in\Omega$ we have $\boldsymbol{i}_\xi(a\otimes_B\omega) = \boldsymbol{i}_\xi(a)\otimes_B\omega$; therefore

$$[\nabla,e^{\boldsymbol{i}_\xi}] = e^{\boldsymbol{i}_\xi}\left(-\boldsymbol{i}_{\delta\xi}\pm\frac{1}{2}[\boldsymbol{i}_\xi,\boldsymbol{i}_{\delta\xi}]\pm\frac{1}{6}[\boldsymbol{i}_\xi,[\boldsymbol{i}_\xi,\boldsymbol{i}_{\delta\xi}]]\pm\cdots\right) = -e^{\boldsymbol{i}_\xi}\boldsymbol{i}_{\delta\xi}.$$

Now it is easy to conclude the proof: we have seen in the proof of Theorem 12.3 that $F_\xi^p = e^{\boldsymbol{i}_\xi}(F^p\otimes B)$ and then every $v\in F_\xi^p\otimes_B\Omega$ can be written as $v = e^{\boldsymbol{i}_\xi}(u)$, with $u\in(F^p\otimes B)\otimes_B\Omega$. Therefore

$$\begin{aligned}\nabla(v) = \nabla(e^{\boldsymbol{i}_\xi}(u)) &= [\nabla,e^{\boldsymbol{i}_\xi}](u) + e^{\boldsymbol{i}_\xi}(\nabla u)\\ &= e^{\boldsymbol{i}_\xi}(-\boldsymbol{i}_{\delta\xi}(u)+\nabla(u))\in e^{\boldsymbol{i}_\xi}((F^{p-1}\otimes B)\otimes_B\Omega) = F_\xi^{p-1}\otimes_B\Omega.\end{aligned}$$

□

References

[1] R. Abraham, J.E. Marsden: *Foundations of mechanics.* Second edition, Benjamin/Cummings Publishing Co., Inc., Advanced Book Program, Reading, Mass. (1978).

[2] F. Catanese: *Moduli of algebraic surfaces.* Theory of moduli (Montecatini Terme, 1985), 1-83, Lecture Notes in Math. **1337**, Springer (1988).

[3] H. Clemens: *Geometry of formal Kuranishi theory.* Adv. Math. **198** (2005), no. 1, 311-365.

[4] B. Fantechi, M. Manetti: *Obstruction calculus for functors of Artin rings, I.* J. Algebra **202** (1998) 541-576.

[5] D. Fiorenza, M. Manetti: *L_∞ structures on mapping cones.* Preprint `arXiv:math.QA/0601312`.

[6] K. Fukaya: *Deformation theory, homological algebra and mirror symmetry.* Geometry and physics of branes (Como, 2001), Ser. High Energy Phys. Cosmol. Gravit., IOP Bristol (2003) 121-209. Electronic version available at `http://www.math.kyoto-u.ac.jp/%7Efukaya/fukaya.html` (may 2006).

[7] W.M. Goldman, J.J. Millson: *The deformation theory of representations of fundamental groups of compact Kähler manifolds.* Inst. Hautes Études Sci. Publ. Math. **67** (1988) 43-96.

[8] W.M. Goldman, J.J. Millson: *The homotopy invariance of the Kuranishi space.* Illinois J. Math. Math. **34** (1990) 337-367.

[9] M. Grassi: *L_∞-algebras and differential graded algebras, coalgebras and Lie algebras.* In: *Seminari di Geometria Algebrica 1998-1999* Scuola Normale Superiore (1999).

[10] M. Green: *Infinitesimal methods in Hodge theory.* Algebraic cycles and Hodge theory (Torino, 1993), Lecture Notes in Math. **1594**, Springer (1994) 1-92.

[11] Ph. Griffiths: *Periods of integrals on algebraic manifolds. II. Local study of the period mapping.* Amer. J. Math. **90** (1968) 805-865.

[12] D. Iacono: Thesis; in preparation.

[13] K. Kodaira: *Complex manifold and deformation of complex structures.* Springer-Verlag (1986).

[14] M. Kontsevich: *Deformation quantization of Poisson manifolds, I.* Letters in Mathematical Physics **66** (2003) 157-216; `arXiv:q-alg/9709040`.

[15] M. Kontsevich, Y. Soibelman: *Deformations of algebras over operads and Deligne's conjecture.* In: G. Dito and D. Sternheimer (eds) *Conférence Moshé Flato 1999, Vol. I (Dijon 1999)*, Kluwer Acad. Publ., Dordrecht (2000) 255-307; `arXiv:math.QA/0001151`.

[16] M. Kontsevich, Y. Soibelman: *Homological mirror symmetry and torus fibrations.* K. Fukaya, (ed.) et al., Symplectic geometry and mirror symmetry. Proceedings of the 4th KIAS annual international conference, Seoul, South Korea, August 14-18, 2000. Singapore: World Scientific. (2001) 203-263; `arXiv:math.SG/0011041`.

[17] T. Lada, M. Markl: *Strongly homotopy Lie algebras.* Comm. Algebra **23** (1995) 2147-2161; `arXiv:hep-th/9406095`.

[18] T. Lada, J. Stasheff: *Introduction to sh Lie algebras for physicists.* Int. J. Theor. Phys. **32** (1993) 1087-1104; `arXiv:hep-th/9209099`.

[19] M. Manetti: *Deformation theory via differential graded Lie algebras.* In *Seminari di Geometria Algebrica 1998-1999* Scuola Normale Superiore (1999).

[20] M. Manetti: *Extended deformation functors.* Int. Math. Res. Not. **14** (2002) 719-756; `arXiv:math.AG/9910071`.

[21] M. Manetti: *Cohomological constraint to deformations of compact Kähler manifolds.* Adv. Math. **186** (2004) 125-142; `arXiv:math.AG/0105175`.

[22] M. Manetti: *Lectures on deformations on complex manifolds.* Rend. Mat. Appl. (7) **24** (2004) 1-183; `arXiv:math.AG/0507286`.

[23] M. Manetti: *Lie description of higher obstructions to deforming submanifolds.* Preprint `arXiv:math.AG/0507287` v2 7Oct2005.

[24] A. Newlander, L. Nirenberg: *Complex analytic coordinates in almost complex manifolds.* Ann. of Math. (2) **65** (1957) 391-404.

[25] D. Quillen: *Rational homotopy theory.* Ann. of Math. **90** (1969) 205-295.

[26] Z. Ran: *Universal variations of Hodge structure and Calabi-Yau-Schottky relations.* Invent. Math. **138** (1999) 425-449.

[27] Z. Ran: *Lie atoms and their deformations.* Preprint `arXiv:math.AG/04122204` v3 11Dec2005.

[28] R. Remmert: *Theory of complex functions.* Springer-Verlag GTM **122** (1991).

[29] D.S. Rim: *Formal deformation theory.* In *SGA 7 I*, Exp. VI. Lecture Notes in Mathematics, **288** Springer-Verlag (1972) 32-132.

[30] M. Schlessinger: *Functors of Artin rings.* Trans. Amer. Math. Soc. **130** (1968) 208-222.

[31] M. Schlessinger, J. Stasheff: *Deformation Theory and Rational Homotopy Type.* Preprint (1979).

[32] F. Schuhmacher: *Deformation of L_∞-algebras.* Preprint `arXiv:math.QA/0405485`.

[33] C. Voisin: *Théorie de Hodge et géométrie algébrique complexe I, II.* Société Mathématique de France, Paris (2002).

Dipartimento di Matematica "Guido Castelnuovo",
Università di Roma "La Sapienza",
P.le Aldo Moro 5, I-00185 Roma Italy.

E-mail address: `fiorenza@mat.uniroma1.it`
URL: `www.mat.uniroma1.it/~fiorenza/`

E-mail address: `manetti@mat.uniroma1.it`
URL: `www.mat.uniroma1.it/people/manetti/`